

TURF - GRAY LEAF SPOT

General Information

GENERAL INFORMATION

Shake well before using. Apply TEE-OFF 4.5F with ground equipment, using sufficient volume of spray to provide thorough coverage. Add required amount of TEE-OFF 4.5F to partially filled tank agitated by mechanical or hydraulic means and then add remaining required amount of water. Continuous agitation is required to keep the material in suspension. Do not tank mix with copper-containing materials or with highly alkaline pesticides, such as Bordeaux mixture or lime sulfur. No claim of compatibility with other pesticides is implied. Use the higher rates under conditions of severe disease pressure. Also, see local State Extension Service recommendations for application schedules. Chemigation instructions follow Directions for Use. Do not apply through any irrigation system unless these instructions are followed.

Not for homeowner use. For use only by certified applicators or those under their immediate supervision. Do not apply with fixed wing or rotary aircraft. Not for use on turf being grown for sale or other commercial use as sod. Do not apply to home orchards/backyard fruit trees after fruit set.

IMPORTANT: If, after using TEE-OFF 4.5F as recommended, treatment is not effective, a tolerant strain of fungi may be present (consult your local Advan representative).

Consult your State Agricultural Experiment Station or your State Agricultural Extension Service for advice on the prompt use of some other suitable fungicide.

Limitations, Restrictions, and Exceptions

TURF - GRAY LEAF SPOT

Apply TEE-OFF 4.5F at 3 1/2 to 5 fluid oz./1,000 sq. ft. beginning when conditions are favorable for disease development. Continue at 14-day intervals. Spray should dry on leaves with no "watering in."

For use on all fine turf applications such as industrial lawns, athletic fields, residential and public areas (home lawns, parks, athletic fields, schools and daycare centers, cemeteries, golf course greens, tees, and fairways of Bentgrasses, Bluegrasses, Bermudagrasses, Fescues, Ryegrasses, St. Augustine grasses or their mixtures. TEE-OFF 4.5F is not phytotoxic to any of the above mentioned grasses when used in accordance with the label. TEE-OFF 4.5F is to be used for the prevention and control of the below mentioned diseases (provides both curative and protective action). Apply recommended amounts in sufficient water to obtain thorough coverage, usually three to five gallons spray solution per 1,000 square feet of turf area. When treating golf greens, always treat aprons and approaches. Use the highest recommended rate under conditions of severe disease pressure. Do not apply with fixed-wing or rotary aircraft. Do not graze animals on treated turf. Do not feed clippings to livestock or poultry.

Residential Public Areas

Maximum Individual Application Rate: 77.5 fl. oz./acre or 1.8 fl. oz./1000 square feet

Minimum Retreatment Interval in Days: 14 Days

Maximum Seasonal Total TEE-OFF 4.5F

Product: 310 fl. oz./acre or 7.12 fl.oz./1000 square feet

Active Ingredient: 10.88 LB AI/acre

Golf Course Greens/Tees/Aprons

Maximum Individual Application Rate: 233 fl. oz./acre or 5.35 fl. oz./1000 square feet

Minimum Retreatment Interval in Days: 14 Days

Maximum Seasonal Total

Product: 620 fl. oz./acre or 14.23 fl.oz./1000 square feet

Active Ingredient: 21.8 LB AI/acre

Golf Course Fairways Outside of Florida

Maximum Individual Application Rate: 155 fl. oz./acre or 3.56 fl. oz./1000 square feet

Minimum Retreatment Interval in Days: 14 Days

Maximum Seasonal Total

Product: 155 fl. oz./acre or 3.56 fl.oz./1000 square feet

Active Ingredient: 5.45 LB AI/acre

Florida Golf Course Fairways Only During Overseeding

Maximum Individual Application Rate: 77.5 fl. oz./acre or 1.8 fl. oz./1000 square feet

Minimum Retreatment Interval in Days: 14 Days

Maximum Seasonal Total

Product: 77.5 fl. oz./acre or 1.8 fl.oz./1000 square feet

Active Ingredient: 2.72 LB AI/acre

Method

[Broadcast/Foliar Ground](#)

Rates

[field_rates 0](#)

-

Restricted Entry Interval

12 hours

Exemption: If the product is applied by drenching, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

Timings

[When conditions are favorable for disease development.](#)