

ORNAMENTALS - ROSES - BLACK SPOT

General Information

GENERAL INFORMATION

Shake well before using. Apply TEE-OFF 4.5F with ground equipment, using sufficient volume of spray to provide thorough coverage. Add required amount of TEE-OFF 4.5F to partially filled tank agitated by mechanical or hydraulic means and then add remaining required amount of water. Continuous agitation is required to keep the material in suspension. Do not tank mix with copper-containing materials or with highly alkaline pesticides, such as Bordeaux mixture or lime sulfur. No claim of compatibility with other pesticides is implied. Use the higher rates under conditions of severe disease pressure. Also, see local State Extension Service recommendations for application schedules. Chemigation instructions follow Directions for Use. Do not apply through any irrigation system unless these instructions are followed.

Not for homeowner use. For use only by certified applicators or those under their immediate supervision. Do not apply with fixed wing or rotary aircraft. Not for use on turf being grown for sale or other commercial use as sod. Do not apply to home orchards/backyard fruit trees after fruit set.

IMPORTANT: If, after using TEE-OFF 4.5F as recommended, treatment is not effective, a tolerant strain of fungi may be present (consult your local Advan representative).

Consult your State Agricultural Experiment Station or your State Agricultural Extension Service for advice on the prompt use of some other suitable fungicide.

Limitations, Restrictions, and Exceptions

ORNAMENTALS - ROSES - BLACK SPOT

Apply early summer or at first sign of disease. Repeat application every 7-14 days as needed during disease period.

TEE-OFF 4.5F is a broad-spectrum systemic fungicide, which controls a variety of

diseases on woody and herbaceous ornamental crops. TEE-OFF 4.5F is also effective as a preplant dip on cutting and bulbs. TEE-OFF 4.5F may be used to control the listed diseases on fruit and nut trees, namely almond, apple, apricot, cherry, nectarine, peach, pecan, plum, and prune trees. Do not use fruit or nuts from trees treated with this product for food or feed.

FOLIAR SPRAY-HYDRAULIC APPLICATION

Mixing instructions: Add required amount of TEE-OFF 4.5F to partially filled tank agitated by mechanical or hydraulic means and then add remaining required amount of water. Maintain continuous agitation to keep the material in suspension and apply with properly calibrated spray equipment. Apply until leaves glisten, but not to point of runoff.

Application Rates and Timing for Diseases: Begin applications when disease first appears and repeat at 10- to 14-day intervals, during the growing season. Shorten the interval during humid, rainy weather.

Ground Application: Use 20 Fluid ounces of TEE-OFF 4.5F per 100 gallons of water. For control of Powdery Mildew and Botrytis (Gray Mold), use 10 fluid ounces of TEE-OFF 4.5F per 100 gallons of water, or 1 teaspoonful per 2 gallons of water.

Method

[Broadcast/Foliar Ground](#)

Rates

[field rates 0](#)

-

Restricted Entry Interval

12 hours

Exemption: If the product is applied by drenching, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

Timings

[Early summer or at first sign of disease.](#)