

DRENCH APPLICATION

General Information

GENERAL INFORMATION

Shake well before using. Apply TEE-OFF 4.5F with ground equipment, using sufficient volume of spray to provide thorough coverage. Add required amount of TEE-OFF 4.5F to partially filled tank agitated by mechanical or hydraulic means and then add remaining required amount of water. Continuous agitation is required to keep the material in suspension. Do not tank mix with copper-containing materials or with highly alkaline pesticides, such as Bordeaux mixture or lime sulfur. No claim of compatibility with other pesticides is implied. Use the higher rates under conditions of severe disease pressure. Also, see local State Extension Service recommendations for application schedules. Chemigation instructions follow Directions for Use. Do not apply through any irrigation system unless these instructions are followed.

Not for homeowner use. For use only by certified applicators or those under their immediate supervision. Do not apply with fixed wing or rotary aircraft. Not for use on turf being grown for sale or other commercial use as sod. Do not apply to home orchards/backyard fruit trees after fruit set.

IMPORTANT: If, after using TEE-OFF 4.5F as recommended, treatment is not effective, a tolerant strain of fungi may be present (consult your local Advan representative).

Consult your State Agricultural Experiment Station or your State Agricultural Extension Service for advice on the prompt use of some other suitable fungicide.

Limitations, Restrictions, and Exceptions

DRENCH APPLICATION

Botrytis: herbaceous annuals, perennials and bedding plants

Fusarium: herbaceous annuals, perennials and bedding plants

Sclerotinia: stem, herbaceous annuals, perennials and crown & roots bedding plants

Cylindrocladium: rot woody ornamentals (such as azaleas, rhododendrons, conifers, poinsettias)

Thielaviopsis: rot woody ornamentals (such as azaleas, rhododendrons, conifers, poinsettias)

NOTE: TEE-OFF 4.5F does not control Pythium spp. or Phytophthora spp.

Application Rates and Timing for Disease Control: Use 10-20 fluid ounces of TEE-OFF 4.5F per 100 gallons of water. Apply as a drench or heavy spray to 800 sq. ft. of bench area (1 to 2 pints per square foot) after transplanting into propagation beds or containers. Repeat at 2- to 4-week intervals during periods favorable for disease.

Refer in the label regarding application rates for specific types of containers.

Method

[Drench](#)

Rates

[field_rates 0](#)

-

Restricted Entry Interval

12 hours

Exemption: If the product is applied by drenching, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

Timings

[After transplanting.](#)