

# **LEAFY VEGETABLES SUCH AS BROCCOLI, CAULIFLOWER, LETTUCE, SPINACH AND CELERY**

## General Information

### GENERAL INFORMATION

CALMAX® PREMIUM contains protein hydrolysate obtained from hydrolysis of protein to amino acids and short polypeptides that are a source of nitrogen. It is designed for foliar application and may be applied alone or in tank mixes containing pesticides. However, when use of an unfamiliar mix is made, a compatibility test is always recommended. Foliar fertilization is a supplement to a regular fertilizer program and will not supply the total nutrients required by a crop. CALMAX PREMIUM is recommended as a foliar nutrition to be applied to crops where disorders due to insufficient calcium are anticipated. This product may also be applied through irrigation systems such as drip or micro sprinklers.

### DIRECTIONS FOR USE

Dissolve the desired amount of CALMAX PREMIUM in water before adding to the spray tank. With proper agitation, CALMAX PREMIUM can be added into the spray tank with agitation running. Do not store solution mixed with pesticide! Spray immediately after mixing. Add CALMAX PREMIUM to the spray tank before adding pesticides.

For aerial application use a maximum of three (3) quarts of CALMAX PREMIUM per 10 gallons of water.

Rates recommended are for standard spray volumes of 10-150 gal/acre. Suggested aerial spray volumes are 5 to 20 gallons/acre. For higher volume sprays, rate should be adjusted. Caution: Store CALMAX PREMIUM in temperatures between 41° F to 104° F and avoid extreme variations in temperature. A reversible separation of ingredients may occur after long storage, this however does not affect quality or effectiveness of the product.

Limitations, Restrictions, and Exceptions

## NUMBER AND TIMING OF APPLICATIONS

Make first application 4-6 leaf stage or one week after transplanting. Make two additional applications at 14 day intervals.

Method

[Spray](#)

Rates

[field\\_rates 0](#)

•

Timings

[First application 4-6 leaf stage or one week after transplanting.](#)