

TREE NUT CROPS (PECAN) - FIRE ANTS, WALNUT HUSK FLY

General Information

Resistance. Some insects are known to develop resistance to products used repeatedly for control. Because the development of resistance cannot be predicted, the use of this product should conform to resistance management strategies established for the use area. Consult your local or state agricultural authorities or universities for details.

If resistance to this product develops in your area, this product, or other products with a similar mode of action, may not provide adequate control. If poor performance cannot be attributed to improper application or extreme weather conditions, a resistant strain of insect pest may be present. If you experience difficulty with control and resistance is a reasonable cause, immediately consult your local company representative or agricultural advisor for the best alternative method of control for your area.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

GENERAL INSTRUCTIONS

Unless otherwise directed by registered supplemental labeling, follow the Directions for Use in each crop group section.

Bifenture 10DF Insecticide/Miticide must be diluted with water before spray application. Do not use nozzle screens (e.g. strainers) greater than 50 mesh size.

Fill the spray tank 1/2 full with water. Add required amount of Bifenture 10DF. Start agitation in the tank. Air agitation is not recommended. Mix thoroughly to fully disperse and suspend the wettable powder. Fill the spray tank with the required amount of water. Maintain agitation during storage in nurse or supply tank and during application.

Mix as needed; do not store diluted material.

Application Instructions

Rate of application is variable according to pest pressure, timing of sprays, and field scouting. Use lower rates under light to moderate infestation; higher rates under heavy insect pressure and for mite control. Arid climates generally require higher rates.

In New York State this product may not be applied within 100 feet (using ground equipment) to 300 feet (using aerial equipment) of coastal marshes or streams that drain into coastal marshes.

Limitations, Restrictions, and Exceptions

TREE NUTS CROPS

- Minimum Spray Intervals: Apply Bifenture 10DF as needed to maintain control, but do not apply at intervals sooner than 15 days.
- Do not graze livestock in treated orchards or cut treated cover crops for feed.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

21 days

Rates

[field rates 0](#)

[field rates 1](#)

•

Restricted Entry Interval

12 hours

Timings

[N.A.](#)