

# **CONIFERS - RHIZOSPHAERA NEEDLECAST, SCHIRRHIA BROWN SPOT**

## General Information

### General Precautions and Restrictions:

- DO NOT apply this product in a way that will contact workers or other persons, or pets, either directly or through drift. Only protected handlers may be in the area during applications. For any requirements specific to your State or Tribe, consult the Agency responsible for pesticide regulation.

### Limitations, Restrictions, and Exceptions

## CONIFERS

Apply this product in sufficient water and with proper calibration to obtain uniform coverage of tree canopy. Application with ground equipment is preferable to aerial application because ground applications generally give better coverage of the tree canopy.

DO NOT allow livestock to graze in treated areas.

DO NOT apply ArmorTech CLT 720 FL within one week before or after application of oil or an oil-based pesticide.

## APPLICATION DIRECTIONS

- Make the first application in spring when new shoot growth is ½ to 2 inches in length. Make additional applications at 3 to 4 week intervals until conditions no longer favor disease development. For use in nursery beds, apply the highest rate specified on a 3 week schedule.

SPRAY VOLUME (GALLONS/ACRE) : 5 to 10 (concentrate ground or aircraft) to 100 (dilute)

### Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field\\_rates 0](#)

[field\\_rates 1](#)

•

Restricted Entry Interval

12 hours

Timings

[When new shoot growth is ½ to 2 inches in length.](#)