

FLOWERING PLANTS AND BULBS - FLOWER SPOTS AND BLIGHTS

General Information

General Precautions and Restrictions:

- DO NOT apply this product in a way that will contact workers or other persons, or pets, either directly or through drift. Only protected handlers may be in the area during applications. For any requirements specific to your State or Tribe, consult the Agency responsible for pesticide regulation.

Limitations, Restrictions, and Exceptions

ORNAMENTAL PLANTS

Apply ArmorTech CLT 720 FL at a rate of 1 3/8 pints per 100 gallons of water unless other directions are given in the tables below. Apply enough diluted spray per acre to provide thorough coverage of all plant parts that are intended to be protected from disease, generally ranging from 20 to 150 gallons per acre. Repeat applications at 7 to 14 day intervals until conditions are no longer favorable for disease. During periods when conditions favor severe disease incidence, generally cloudy or wet weather, apply ArmorTech CLT 720 FL at 7 day intervals. DO NOT apply more than a total of 36.4 lbs. chlorothalonil active ingredient per acre per growing season on field-grown ornamentals.

Fruits and other structures which may be borne on treated plants MUST NOT BE EATEN.

Not approved for use in California: Arabian violet

Method

[N.A.](#)

Rates

[field rates 0](#)

-

Restricted Entry Interval

12 hours

Timings

N.A.