

RANGELAND, ESTABLISHED GRASS PASTURES (INCLUDING PERENNIAL GRASSLANDS NOT IN AGRICULTURAL PRODUCTION SUCH AS CONSERVATION RESERVE PROGRAM ACRES) - SUSCEPTIBLE ANNUAL AND BIENNIAL BROADLEAF WEEDS

General Information

General Information:

Generally, the lower dosages given will be satisfactory for young, succulent growth of sensitive weed species. For less sensitive species and under conditions where control is more difficult, the higher dosages will be needed. Apply Cornbelt 4 lb. Amine during warm weather when weeds are young and growing actively. Use enough spray volume for uniform coverage by ground or air application. If only bands or rows are treated, leaving middles unsprayed, the dosage per crop acre is reduced proportionately. Read and follow all Use Precautions given on the label.

DO NOT USE THIS PRODUCT ON CROPS GROWN FOR SEED. SOME INBREDS HAVE A VERY LOW TOLERANCE TO CHEMICALS AND THE USE OF THIS COMPOUND COULD RESULT IN INJURY TO A CROP GROWN FOR SEED.

To Prepare the Spray: Mix Cornbelt 4 lb. Amine only with water, unless otherwise directed on the label. Add about half the water to the mixing tank, then add the Cornbelt 4 lb. Amine Herbicide with agitation and finally the rest of the water with continuing agitation. Note: Adding oil, wetting agent, or other surfactant to the spray may increase effectiveness on weeds, but also may reduce selectivity to crops resulting in crop damage.

Consult your State Agricultural Experiment Stations or Extension Service Weed Specialists in many states for recommendations from the label that best fit local conditions. Be sure that use of this product conforms to all applicable regulations.

Apply this product only as specified on the label.

USE PRECAUTIONS

Do not apply this product through any type of irrigation system.

Excessive amounts of 2,4-D in the soil may temporarily inhibit seed germination or plant growth.

Susceptible Plants

Do not apply under circumstances where spray drift may occur to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption. Susceptible crops include, but are not limited to cotton, okra, flowers, grapes (in growing stage), fruit trees (foliage), soybeans (vegetative stage), ornamentals, sunflowers, tomatoes, beans, and other vegetables, or tobacco. Small amounts of spray drift that might not be visible may injure susceptible broadleaf plants.

Limitations, Restrictions, and Exceptions

RANGELAND, ESTABLISHED GRASS PASTURES (INCLUDING PERENNIAL GRASSLANDS NOT IN AGRICULTURAL PRODUCTION SUCH AS CONSERVATION RESERVE PROGRAM ACRES):

Restrictions:

- Do not use on bentgrass, alfalfa, clover or other legumes.
- Do not use on newly seeded areas until grass is well established.
- Do not use from early boot to milk stage where grass seed production is desired.
- Do not make more than 2 applications per season.
- Do not apply within 30 days of a previous application
- Do not harvest forage for hay within 7 days of application.
- For grazed areas, the maximum use rate is 4.21 pt/acre of Cornbelt 4 lb Amine (2.0 lb of acid equivalent) per application.
- If grass is to be cut for hay, Agricultural Use Requirements for the Worker Protection Standard are applicable.
- Do not apply more than 8.42 pts/acre (4.0 lbs of acid equivalent) of Cornbelt 4 lb

Amine per use season

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

7 days

Rates

[field_rates 0](#)

[field_rates 1](#)

•

Restricted Entry Interval

12 hours

Timings

[N.A.](#)