

## **ESTABLISHED BEDS (OVER ONE YEAR OLD) : NORTHEAST - IN BEDS WITH SANDY SOILS**

### General Information

#### GENERAL USE PRECAUTIONS

- Read all label directions before using.
- Use only for at specified rates. Do not overdose.
- Do not apply when weather conditions favor drift from areas treated.
- Do not contaminate irrigation water or water used for domestic purposes.
- DEVRINOL 10-G herbicide may be used on mineral soils only. Do not use on soils with over 10% organic matter.
- Treating ground heavily covered with leaves or trash may adversely affect weed control.
- Do not apply to frozen ground.
- To avoid injury to crops not specified on the label, do not plant until 12 months after the last DEVRINOL 10-G herbicide application. A deep disc plow or moldboard plow operations must be carried out before planting.

#### CROP USE DIRECTIONS

Apply DEVRINOL 10-G to a weed free soil surface at rates specified in the following crop use directions sections. Apply in the fall through early spring prior to weed emergence. For best results, the treatment must reach the zone of weed seed germination. If rainfall does not occur, the treatment must be shallowly incorporated or irrigated-in (sprinkler or flood) following application with sufficient water to wet the soil to a depth of 2 to 4 inches. In the Western and Pacific Northwest Regions treatments made in fall through early spring must be incorporated or irrigated-in if no rainfall occurs within two weeks of application. For treatments made other than in fall through early spring in these Regions, and for all treatments applied in the Northern, Southwestern and Southeastern Regions

regardless of season, the treatment must be incorporated or irrigated-in within 24 hours if no rainfall occurs.

Avoid contact with fruit or foliage. Do not apply when fruit or nuts are on the ground during the harvest period. Do not graze treated areas. Application may be made immediately after planting, except where treatment is restricted to established crops only. For these uses, apply to stands that have been established in the field for at least one year.

NOTE: See Crop Use Directions Section for weeds controlled in Cranberries.

NOTE: DEVRINOL 10-G does not control established weeds.

### Limitations, Restrictions, and Exceptions

#### Established beds (over one year old)

For control of nutsedge (*Cyperus dentatus*) and rice cutgrass (*Leersia oryzoides*) and purplestem beggarticks (*Bidens connata*), apply DEVRINOL 10-G at the rates indicated below to a weed free soil surface before spring growth begins or in the fall after harvest. Do not apply when beds are under winter flood.

#### Method

[Surface](#)

[Spray](#)

#### Rates

[field rates 0](#)

- 

#### Restricted Entry Interval

24 hours

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

#### Soils

[Sand](#)

#### Timings

[Before spring growth begins or in the fall after harvest.](#)