

PEANUTS - BEAN LEAF BEETLE ETC.

General Information

GENERAL DIRECTIONS FOR USE

Thorough crop coverage is necessary for both initial and residual control. Apply by ground in at least 10 gals./A or by air in at least 2 gals./A using sufficient water to obtain full coverage of foliage unless the label specifies otherwise. In situations where foliage is dense or pest pressure is high (heavier insect or egg pressure, larger larval stages), control can be improved by use of higher application volumes and/or higher use rates.

For cutworm control, LAMBDA-CY EC INSECTICIDE-RUP may be applied before, during, or after planting. When making soil incorporated applications, use higher rates for better control.

Resistance Management

LAMBDA-CY EC INSECTICIDE-RUP is a Group 3 Insecticide (contains the active ingredient lambda-cyhalothrin). Some insects are known to develop resistance to products used repeatedly for control. Because the development of resistance cannot be predicted, the use of this product should conform to resistance management strategies established for the use area. Consult your local or state agricultural authorities for details.

If resistance to this product develops in your area, this product, or other products with a similar mode of action, may not provide adequate control. If poor performance cannot be attributed to improper application or extreme weather conditions, a resistant strain of insect may be present. If you experience difficulty with control and resistance is a reasonable cause, immediately consult your local company representative or agricultural advisor for the best alternative method of control for your area.

Limitations, Restrictions, and Exceptions

PEANUTS - BEAN LEAF BEETLE ETC.

Use scouting to determine need for application, usually at intervals of 5 or more days. Base the timing and frequency of applications on when insect populations reach local economic thresholds.

Do not graze or harvest treated soybean forage, straw or hay for livestock feed.

Note:

Fall Armyworm: Use higher rates for large larvae.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

30 days

Rates

[field_rates 0](#)

[field_rates 1](#)

•

Restricted Entry Interval

24 hours

Timings

[When insect populations reach local economic thresholds.](#)