

## **LAWN - (0.1 - 0.2 LBS. PER ACRE)**

### General Information

### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply more than 0.2 lb. active ingredient (a.i.) per acre per application on residential use sites (i.e., around private homes, apartment buildings, condominiums, non-agricultural outbuildings, non-commercial greenhouses, pre-schools or day care facilities). Application at up to 0.4 lb. a.i. per acre per application is allowed on non-residential use sites (i.e., around institutional, public, commercial or industrial buildings; parks; recreational areas or athletic fields).

Do not broadcast apply more than 0.4 lbs. a.i. (200 lbs. UP-Star Easy Spread Granular) per acre per year.

For ground application only. Do not apply by air.

Do not apply within 25 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish farm ponds.

Do not use on golf courses, sod farms, nurseries, in commercial greenhouses or on grass grown for seed.

### Limitations, Restrictions, and Exceptions

Active Ingredient lbs. per acre: 0.1-0.2 lbs. per acre

Rate UP-Star Easy Spread Granular: 50-100 lbs./A Or 1.15-2.3 lbs./ 1,000 sq. ft.

Annual Bluegrass Weevil (*Hyperodes*) Adults:

Time application to control adult weevils as they leave their over-wintering sites and move into grass areas. This movement usually starts when Forsythia is in full bloom and finishes when flowering dogwood is in full bloom. Consult your State Cooperative Extension Service for more specific information regarding application

timing.

#### Billbug Adults:

Make first application when adult billbugs are first seen during April and May. In temperate regions, spring applications for billbug adults will also control overwintered chinch bugs.

#### Black Turfgrass Ataenius Adults:

Apply in May and July to control the first and second generation of black turfgrass ataenius adults, respectively. Time the May application to coincide with the full bloom stage of Vanhoutte spiraea (*Spiraea vanhouttei*) and horse chestnut (*Aesculus hippocastanum*). Time the July application to match the blooming of Rose of Sharon (*Hibiscus syriacus*).

#### Method

##### [Broadcast](#)

##### Rates

##### [field rates 0](#)

##### [field rates 1](#)

##### [field rates 2](#)

- 

#### Timings

[Billbug Adults: When adult billbugs are first seen during April and May.](#)

[Black Turfgrass Ataenius : In May and July.](#)