

STRAWBERRIES, RASPBERRIES, BLUEBERRIES

General Information

DIRECTIONS FOR USE

Not for use on plants being grown for sale or for other commercial use, or for commercial seed production or for research purposes.

Step #1: For vegetables, measure the area to be sprayed. Calculate square feet by multiplying length by width. For all other applications, apply as a thorough cover spray.

Step #2: Use a clean sprayer. Carefully measure and mix the amount of concentrate and water as directed.

Step #3: Spray as directed. Thorough coverage is important. Flush sprayer with clean water after each use.

Make applications when insects first appear. Repeat applications as directed. See directions for complete insect list.

People and pets should not be allowed in treated areas until spray has dried.

GENERAL DIRECTIONS

Use 2 fl. oz. in 1 gal. of water. Apply when insects appear. Wet the plants to the dripping point. Try to hit underside of leaves and penetrate dense foliage.

Limitations, Restrictions, and Exceptions

VEGETABLES

For all of the following vegetables, apply 1 gallon of finished mix to 1,000 sq. ft.

How to use:

Use 1 ½ fl. oz. in 1 gal. Allow 14 days between last application and harvest. Make no more than 8 foliar applications per growing season.

Method

[Foliar spray](#)

Rates

[field rates 0](#)

[field rates 1](#)

-

Timings

[N. A.](#)