

## **FOR CONTROL OF EARWIGS, CUTWORMS, ETC.**

### General Information

- For use around vegetables, fruit trees, berries and gardens
- Kills cutworms, ants, earwigs, slugs and snails
- Easy to use pellet formulation
- Contains spinosad
- Lures slugs, snails and insects from hiding places and plants
- No slime trails
- Lasts up to 4 weeks

Apply to ants excluding fire ants, harvester ants, Pharaoh's ants and carpenter ants.

### APPLICATION DIRECTIONS

Scatter the bait granules on the soil around or near the plants to be protected with a suitable hand or power operated spreader (gravity or rotary) to ensure uniform coverage over the treatment area. Apply bait evenly at approximately 0.5 to 1 lb per 1000 square feet, or for smaller areas, 0.075 to 0.15 oz, or 0.5 to 1 teaspoon, per square yard. Do not place in piles. Scatter the bait around the perimeter of the area to provide a protective "barrier" for slugs and snails entering the area. If the ground is dry, wet it before applying bait. The product works best when soil is moist but with little or no standing water. Do not use food utensils such as tablespoons, cups, etc. for food purposes after pesticide use.

Do not apply more than 3 times in any 30 day period. Apply at the higher rate if the infestation is severe. Re-apply after heavy rain or watering. Do not apply within 3 days of harvest.

For seedlings spread the bait around the base of the stem. For trees or shrubs, spread the bait around the base of the tree. For small plots, treat around the base of the plants to be protected. In greenhouses, scatter the bait in the pots of plants being damaged or around pots on greenhouse benches.

Evening is the best time to apply the bait as pests travel and feed mostly by night or early morning. Treat all likely areas of infestation, especially around the perimeter of garden plots, for target pests because pests travel into plant areas

from daytime refuges.

## Limitations, Restrictions, and Exceptions

### FOR CONTROL OF EARWIGS, CUTWORMS, ETC.

#### APPLICATION DIRECTIONS

Scatter the bait granules on the soil around or near the plants to be protected with a suitable hand or power operated spreader (gravity or rotary) to ensure uniform coverage over the treatment area. Apply bait evenly at approximately 0.5 to 1 lb per 1000 square feet, or for smaller areas, 0.075 to 0.15 oz, or 0.5 to 1 teaspoon, per square yard. Do not place in piles. For slugs and snails also scatter the bait around the perimeter of the area to provide a protective \"barrier\" for slugs and snails entering the area. If the ground is dry, wet it before applying bait. The product works best when soil is moist but with little or no standing water. Do not use food utensils such as tablespoons, cups, etc. for food purposes after pesticide use.

Do not apply more than 3 times in any 30 day period. Apply at the higher rate if the infestation is severe. Re-apply after heavy rain or watering.

For seedlings spread the bait around the base of the stem. For trees or shrubs, spread the bait around the base of the tree. For small plots, treat around the base of the plants to be protected. In greenhouses, scatter the bait in the pots of plants being damaged or around pots on greenhouse benches.

Apply to ants excluding fire ants, harvester ants, Pharaoh's ants and carpenter ants.

#### Method

##### [Bait](#)

##### Rates

##### [field\\_rates 0](#)

##### [field\\_rates 1](#)

##### [field\\_rates 2](#)

- 

#### Timings

[When soil is moist but with little or no standing water.](#)