

## **VEGETABLE DISEASES - CUCUMBERS ETC.**

### General Information

#### GENERAL INFORMATION

CONTROLS: Leaf spots, rust, blights, fruit rots, mildews, scab, molds, etc.

USE ON: vegetables, fruit trees, flowers, shrubs, and shade trees

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in interior plantscapes, ornamental gardens or parks, and grounds. Also for use on fruit trees and vegetables.

**READ ENTIRE LABEL AND USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS.**

This product is highly effective for the control of a broad spectrum of vegetable, fruit and ornamental plant diseases when it is used according to the directions on the label. When conditions are favorable for development, thorough uniform coverage of plant surfaces is essential for good disease control.

General Precautions and Restrictions: Do not allow people or pets to enter treated areas until sprays have dried. Do not apply this product in a way that will contact other persons or pets, either directly or through drift.

Limitations, Restrictions, and Exceptions

VEGETABLE DISEASES

Spray foliage to point of runoff. Repeat at 7 to 14 day intervals. Use the shorter spray schedule during severe disease conditions or wet weather unless otherwise stated.

Note: Avoid spraying plants during extremely hot and sunny weather. Do not apply to wilted plants. Do not graze treated areas or feed treated plant parts to livestock.

CUCUMBERS, MELONS, PUMPKINS, SQUASH: Anthracnose, Downy Mildew and Target Spot; Scab, Gummy Stem Blight (Black Rot) Leaf Blight, Cercospora Leaf Spot, Powdery Mildew (except Southwestern states). Also suppresses Fruit Rot (Rhizoctonia) on cucumbers - Begin when plants are in the first true leaf stage or when conditions are favorable for disease. Repeat applications at 7 day intervals or more frequently if required. Do not apply more than 50 gal. per 1,000 sq. ft. per season.

Method

[Spray](#)

Timings

[When plants are in the first true leaf stage or when conditions are favorable for disease.](#)