

FOR ST. AUGUSTINEGRASS AND BENTGRASS

General Information

DIRECTIONS FOR USE

USE RESTRICTIONS:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift.
- Keep people and pets out of the area during application.
- Do not allow people or pets to enter the treated area until sprays have dried.
- Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Do not spray where grass is emerging from dormancy. Do not apply to newly seeded lawn grasses until well established. Do not spray when the temperature exceeds 85°F. Do not use on carpetgrass, dichondra, centipedegrass, nor on lawns or turf where desirable clovers are present. Do not exceed specified dosages for any area.

- Do not apply as a fine mist because of potential for injury to desirable plants.
- Do not use this product on or near desirable plants, including contact of spray on exposed root systems or adventitious shoots within the drip line of desirable trees and shrubs, since injury may result.

Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. To prevent product run-off, do not over water the treated area to the point of runoff or apply when raining or when rain is expected that day. Rinse applicator over lawn area only.

WHERE TO USE

BONIDE Liquid Weed & Feed may be used on bentgrass, bermudagrass, Kentucky bluegrass, fescues, perennial ryegrass, and St. Augustinegrass lawns, vacant lots, playgrounds where broadleaf weeds are present. Bonide Liquid Weed & Feed controls listed broadleaf weeds, including sensitive weeds such as chickweed, clover, dandelion, knotweed and plantain.

WHEN TO USE

1. Spray during the growing season when weeds are actively growing.
2. On new lawns, wait until the grass has reached a height of 2 inches before making an overall spray treatment. Wait three (3) weeks after application before sowing seed or planting stolons.
3. Poor weed control may result if spray is applied during drought or just before rain.
4. For best results, mow 1 to 2 days before treating.
5. Do not water lawn within 24 hours after treatment.

HOW TO USE

Shake Well Before Using.

For Operation of Hoes-End Sprayer

Make sure knob control lever is in the "OFF" position. shake well and attach to hose. Turn water on at faucet, aim nozzle toward lawn and turn knob control lever to "ON" to begin spraying. When finished, turn control lever to "OFF" position. turn faucet to "OFF" position. discard residual water pressure by turning "ON" and "OFF" again.

Limitations on broadcast applications to turfgrass:

The maximum application rate is 13 fl. oz. of product per 1,000 sq. ft. per application. The Maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 26 fl. oz. of product per 1,000 sq. ft. excluding spot treatments.

Limitations on spot treatments to residential turfgrass:

Spot treatment is defined as a treatment area no greater than 1,000 sq. ft. per acre. The maximum application rate is 13 fl. oz. per 1,000 sq. ft. per application.

Limitations, Restrictions, and Exceptions

FOR ST. AUGUSTINEGRASS AND BENTGRASS

Apply 8 fluid ounces (16 tablespoons) of product per 1,000 square feet of turfgrass. This product at this rate provides 0.13 pounds of nitrogen per 1,000 square feet per application.

Do not apply this product to 'Floritam' St. Augustinegrass in Florida.

Refer in the label for Area to Treat, Square Feet (Sq. Ft.).

Method

[Spot treatment](#)

[Broadcast](#)

[Spot treatment](#)

[Broadcast](#)

Rates

[field_rates 0](#)

[field_rates 1](#)

•

Timings

[Preemergence \(Weed\)](#)

[Postemergence \(Weed\)](#)