

FUNGICIDE USE

General Information

GENERAL INFORMATION

- Three garden products in one.
- Multi-purpose fungicide/insecticide/miticide-3 in 1 product.
- Kills eggs, larvae and adult stages of insects.
- Prevents fungal attack of plant tissues.
- Not for use in food-handling areas.
- Shake well before using.
- Thorough plant coverage is essential to obtain optimal levels of control.
- Do not apply to wilted or otherwise stressed plants, or to newly transplanted material prior to root establishment.
- As with other pest control products, exercise care in timing applications to early morning/late evening to minimize the potential for leaf burn.
- Use with care on plants with tender tissue. Check for leaf burn in small-scale trials prior to use.

Limitations, Restrictions, and Exceptions

FUNGICIDE USE

Application Instructions

- This product is an effective fungicide for the prevention and control of various fungal diseases including black spot on roses, powdery mildew, downy mildew, anthracnose, rust leaf spot, botrytis, needle rust, scab, and flower, twig and tip blight.
- This product may be used on all roses, flowers, houseplants, ornamental trees and shrubs, fruits, nuts, and vegetables grown in and around the home and home garden.
- As a preventative, apply this product on a 7 to 14 day schedule until the potential for disease development is no longer present. Spray to run off. Complete coverage of all plant tissue is necessary for control.
- To control diseases already present, apply this product on a 7- day schedule until disease pressure is eliminated. Then continue spraying on 14-day schedule to

prevent the disease from recurring. Spray to run off. Complete coverage of all plant tissue is necessary for control.

- To prevent rust leaf spot diseases, anthracnose, and scab, begin applications at the first sign of spring budding.

- To prevent powdery mildew, apply in mid-summer or when disease is first detected. Continue applications until disease pressure no longer exists.

- To prevent black spot on roses, begin applications at first bud emergence in spring. Apply this product on a 14-day preventative schedule through the spring and fall to prevent disease emergence.

Method

Spray

Timings

Powdery mildew: In mid-summer or when disease is first detected.

Rust Leaf Spot, anthracnose, scab: at the first sign of spring budding.

Black spot on roses: at first bud emergence in spring.