

# **TOMATO BLOSSOM-END ROT**

## General Information

### HELPFUL INFORMATION:

Tomato blossom-end rot is a disorder caused by calcium deficiency. The disorder can be aggravated by the application of too much nitrogen fertilizer to tomato plants. Blossom-end rot appears as a round, brown sunken area on the end of the tomato fruit opposite to the stem. As a fruit ripens the area turns dry and leathery. The disorder often appears after a period of rapid growth followed by dry conditions, or in periods of heavy rain that caused calcium to leach from the soil.

Helps correct calcium deficiencies.

Also use on cucumbers, melons and peppers.

### Limitations, Restrictions, and Exceptions

**DIRECTIONS FOR USE:** Apply Only As Directed. Apply Bonide Rot-Stop Ready-to-Use undiluted. Apply to the point of run-off onto the foliage and fruit during periods of rapid growth or following excessive rain fall. Apply in early morning or evening when temperatures are lower to avoid foliage burn. Make applications 5 to 7 days apart when conditions are favorable for appearance of the disorder. Regular spray programs are not recommended as prolonged spraying and rates in excess of those specified can cause foliage burn.

### Method

[Spray](#)

### Timings

[Apply in early morning or evening when temperatures are lower to avoid foliage burn.](#)