

PRESSURE OR TANK TYPE SPRAYERS - CENTIPEDEGRASS AND ST. AUGUSTINEGRASS

General Information

USE RESTRICTIONS

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until sprays have dried.
- Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.
- Use only lawn-type sprayers. Do not spray on carpetgrass, dichondra, nor on lawns or turf where desirable clovers are present. Do not exceed specified dosages for any area.
- The turf tolerance of centipedegrass and St. Augustinegrass can range from sensitive to tolerant. Do not apply this product to 'Floritam' St. Augustinegrass in Florida.
- Do not apply as a fine mist because of potential for injury to desirable plants.
- Do not use this product on or near desirable plants, including contact of spray on exposed root systems or adventitious shoots within the drip line of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. To prevent product run-off, do not over water the treated area to the point of runoff or apply when raining or when rain is expected that day. Rinse sprayer over lawn area only.

WHERE TO USE

Use on bahiagrass, bermudagrass, Kentucky bluegrass, fescue, perennial ryegrass, zoysiagrass, centipedegrass, and St. Augustinegrass lawns, vacant lots, playgrounds and other similar areas where broadleaf weeds are present. This product is especially effective on all hard-to-kill weeds such as poison ivy, poison oak, sheep sorrel and spurge. In addition, this product controls most other broadleaf weeds, including sensitive weeds such as chickweed, clover, dandelion, knotweed

and plantain. A list is given on a following panel.

WHEN TO USE

1. Spray during the growing season when weeds are actively growing.
2. On new lawns, wait until the grass has reached a height of 2 inches before making an overall spray treatment. However, a spot treatment may be made at any time with a small pressure sprayer. Wait 3 weeks before seeding or planting stolons, plugs, or sprigs.
3. Poor weed control may result if spray is applied during drought or just before rain.
4. For best results, mow before treating.
5. Do not water lawn within 24 hours after treatment.

LIMITATIONS ON BROADCAST APPLICATIONS TO RESIDENTIAL TURFGRASS

The maximum application rate is 4.0 fl. oz. of product per 1,000 sq. ft. per application. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 8.0 fl. oz. of product per 1,000 sq. ft.

SPOT SPRAYING

To spray small areas or individual weeds, do not use a hose end sprayer. Refer to Chart 3 or Chart 4 for mixing instructions. Spray as needed and be certain to wet the weeds.

LIMITATIONS ON SPOT TREATMENTS TO RESIDENTIAL TURFGRASS

Spot treatment is defined as a treatment area no greater than 1,000 sq. ft. per acre. The maximum application rate is 4.0 fl. oz. per 1,000 sq. ft. per application. The maximum number of spot treatments is limited to 2 per year with a minimum of 30 days between applications.

USE PRECAUTIONS

- Schedule the herbicide applications when air temperatures are 60° to 90°F with good soil moisture. Avoid applications during the spring transition of the warm season grasses and during periods of extremely high temperatures. The tolerance of warm season grasses decreases at air temperatures greater than 90°F and when turfgrass is drought stressed.

Limitations, Restrictions, and Exceptions

PRESSURE SPRAYERS

Measure indicated amount of this product, pour into tank, and dilute with water according to the following chart. Use a spray pattern that gives small spray droplets without any fine mist, and cover the area indicated.

Method

[Spray](#)

Rates

[field_rates 0](#)

[field_rates 1](#)

-

Timings

[When weeds are actively growing.](#)