

LAWNS - BROADLEAF WEEDS CONTROL

General Information

PRECAUTIONS AND RESTRICTIONS:

- Do not apply this product in a way that will contact any person, or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until sprays have dried.
- Not for use on turf being grown for sale or other commercial use on sod, or for commercial seed production, or for research purposes.

PRODUCT FACTS

This product contains four active ingredients including carfentrazone-ethyl that broaden the spectrum of weed control.

This product offers these advantages:

- Excellent postemergent activity with proven performance for broadleaf weed control in turfgrass.
- Superior cool weather performance.
- Rapid and effective weed control for common and troublesome weed species in turfgrass, e.g. spurge, pennywort (dollarweed), dandelion, and white clover.
- Injury to weeds can be noticed within hours of the application and plant death can occur within 7-14 days.

WHERE TO USE

On Lawns:

Use to spot treat broadleaf weeds commonly found in cool season and warm season grasses such as: Kentucky bluegrass, Perennial ryegrass, Tall fescue, Red or fine leaf fescues, Mixtures of cool season species, Common bermudagrass, Hybrid bermudagrass, St. Augustinegrass (excluding 'Floritam'), Zoysiagrass.

Prohibitions:

- Do not apply this product to bentgrass, carpetgrass, dichondra, legumes, and lawns where desirable clovers are present.

- Do not apply to 'Floritam' St. Augustinegrass grown in Florida.
- Do not allow spray to contact vegetables, flowers or ornamentals as injury may result.
- Do not apply more than 1.9 gallons of product per 1,000 sq. ft. per year.
- Do not apply more than 2 applications per year with a minimum retreatment interval of 21 days.
- Do not apply as a fine mist because of potential for injury to desirable plants.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. To prevent product run-off, do not over water the treated area to the point of runoff or apply when raining or when rain is expected that day.

APPLICATION SCHEDULES (WHEN TO APPLY)

For best results, treat weeds in the spring and the fall when weeds are actively growing. However, this product is effective any time during the growing season.

- Do not apply when air temperatures are above 85°F as lawn injury can occur.
- Hard-to-kill weeds may require a repeat application in 3 weeks.
- Do not water lawn within 12 hours of application. If rainfall occurs within 12 hours after application, retreatment may be necessary.
- Do not apply to newly seeded lawns until after the second mowing.
- Delay application to newly sodded, plugged, or sprigged lawns until new grass has been mowed four times.
- Treated areas may be reseeded 2 weeks after application.
- Applications to dormant bermudagrass, zoysiagrass, and St. Augustinegrass may be used to control actively growing broadleaf weeds.

HOW TO USE

For Containers with Trigger Sprayers: This product is applied directly from the container fitted with the trigger sprayer. Shake the container well. Direct the spray to the target weeds to provide uniform coverage of the upper leaf surfaces. Apply to the point of runoff only. Excessive spraying can injure lawns.

Limitations, Restrictions, and Exceptions

WHERE TO USE

On Lawns:

Use to spot treat broadleaf weeds commonly found in cool season and warm season grasses such as: Kentucky bluegrass, Perennial ryegrass, Tall fescue, Red or fine leaf fescues, Mixtures of cool season species, Common bermudagrass, Hybrid bermudagrass, St. Augustinegrass (excluding 'Floritam'), Zoysiagrass.

Prohibitions:

- Do not apply this product to bentgrass, carpetgrass, dichondra, legumes, and lawns where desirable clovers are present.
- Do not apply to 'Floritam' St. Augustinegrass grown in Florida.
- Do not allow spray to contact vegetables, flowers or ornamentals as injury may result.

APPLICATION SCHEDULES

For best results, treat weeds in the spring and the fall when weeds are actively growing. However, this product is effective any time during the growing season.

- Do not apply when air temperatures are above 85°F as lawn injury can occur.
- Hard-to-kill weeds may require a repeat application in 2 to 3 weeks.
- Do not water lawn within 12 hours of application. If rainfall occurs within 12 hours after application, retreatment may be necessary.
- Do not apply to newly seeded lawns until after the second mowing.
- Delay application to newly sodded, plugged, or sprigged lawns until new grass has been mowed four times.
- Treated areas may be reseeded 2 weeks after application.
- Applications to dormant bermudagrass, zoysiagrass, and St. Augustinegrass may be used to control actively growing broadleaf weeds.

- Do not apply more than 1.5 lb ai (MCPA)/acre per year. Do not apply more than 2 applications per year with a minimum retreatment interval of 21 days.

HOW TO USE

For Containers with Trigger Sprayers: This product is applied directly from the container fitted with the trigger sprayer. Shake the container well. Direct the spray to the target weeds to provide uniform coverage of the upper leaf surfaces. Apply to the point of runoff only. Excessive spraying can injure lawns.

This product will control or suppress the broadleaf weeds.

Method

[Spot treatment](#)

[Directed](#)

Timings

[In the spring and the fall when weeds are actively growing.](#)