

ROOTING WOODY AND SEMI-WOODY PLANTS AND PROPAGATE EVERGREENS

Limitations, Restrictions, and Exceptions

INSTRUCTIONS FOR USE:

Do not use or store near food or feed. Do not use on plants that are to be used for food or feed.

1. There should be as little time as possible between the time of making cuttings, treating with rooting powder and planting.
2. Cuttings should be moistened before treatment, then excess moisture shaken off. Do not wet “hairy” cuttings.
3. Dip basal ends of cuttings into Hormex Rooting Powder.
4. Shake off excess powder by tapping cuttings.
5. Plant treated cuttings in rooting media, being careful not to rub off rooting powder when planting.

Inhibition of root formation is possible by the use of too high a concentration of rooting powder. It is essential for successful propagation that the proper strength rooting powder for the species involved be chosen. Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

Method

[Dip](#)

Restricted Entry Interval

12 hours

EXCEPTION: If the product is soil injected or soil incorporated, the Worker Protection

Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

Timings

[N. A.](#)