

MODERATELY TOLERANT TURFGRASSES

General Information

DIRECTIONS FOR USE

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

This product is for use on Ornamental Turf Lawns (Residential, Industrial and Institutional).

USE PRECAUTIONS

Avoid drift of spray mist to vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not pour spray solutions near desirable plants. Avoid fine mists. Use lawn type sprayer with coarse spray as wind drift is less likely. Avoid contact with exposed feeder roots of ornamentals and trees. Maximum control of weeds will be obtained from spring or early fall applications when weeds are actively growing. Use the higher rate for hard-to-control weeds. Do not exceed specified dosages for any area; be particularly careful within the dripline of tree and other ornamental species. Avoid broadcast applications when air temperature exceeds 90°F. When using small, spot treatment applications in temperatures over 90°F, turf injury may occur.

Do not use clippings as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.

The suitable use of this product on non-recommended turf species may be determined by treating a small area at any rate/1,000 sq. ft. which does not exceed the maximum labeled rate for the turf type where the product is to be applied. The treated area should be observed for any sign of turf injury for a period of 30 days of normal growing conditions to determine the phytotoxicity and efficacy to the treated area.

Do not apply this product by air or through any type of irrigation equipment.

Clean and rinse spray equipment using soap or detergent and water, and rinse thoroughly before reuse for other sprays.

Make no more than 2 applications per year. For ornamental lawns and turf, apply no more than 0.5 oz. ae 2,4-D per 1,000 sq. ft. per application and no more than 1.0 oz. ae 2,4-D per 1,000 sq. ft. per year, excluding spot treatments.

This product combines three proven active ingredients [Quinclorac, 2,4-D, and Dicamba] to provide broad spectrum control of grassy and broadleaf weeds. This herbicide is absorbed by foliage and roots of weeds and translocated throughout the plant.

USE TIPS

GENERAL TIPS

- Shake well before using.
- Spray during the growing season when weeds are young and growing actively.
- For best results, do not mow for 2 days before or after treating.
- Avoid entire lawn treatments when air temperature exceeds 90°F.
- Do not allow spray to get on ornamental plants or vegetables since injury may result. Do not spray around roots of ornamentals and trees.
- Do not use on lawns with desirable clovers and legumes or on ornamentals.
- Avoid sprays which are too fine because they can drift and damage other plants.
- If certain tough weeds are not dead within 3 weeks, repeat with spot treatment application.

TIMING

Apply this product to young, actively growing weeds as a postemergence entire lawn or spot spray. Follow-up applications may be required for dense infestations of broadleaf and grassy weeds.

- Applications of this product made to annual grasses at 2 to 4 tiller may not provide complete control.
- Newly Seeded Areas: Application to grass seedlings is not recommended until after the third mowing.
- Newly Sodded, Sprigged or Plugged Areas: Application should be delayed until 3 to 4 weeks after the sodding, sprigging or plugging operations. Delay applications for 4 weeks after seeding and emergence of Kentucky bluegrass, perennial ryegrass and fine fescues.
- Seeding: Delay application for 4 weeks after seeding and emergence of turf species listed on the label.
- Mowing: Do not mow for two days before or two days after application. Clippings for the first three mowings should be left in the application area. Do not use clippings as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.

IRRIGATION AND RAINFALL

- If soil moisture is not sufficient prior to application, irrigation may improve weed control. For best results, do not water or irrigate for 24 hours after application. If rainfall does not occur in 2 to 7 days after application, irrigation of at least one-half inch is required.

Limitations, Restrictions, and Exceptions

Bluegrass, Annual (Poa Annua), Bermudagrass, Common, Bermudagrass (Hybrids) : Temporary yellowing and discoloration on these species may occur after application.

Fescue, Fine: Only apply to fine fescue when it is part of a blend.

HOW TO USE:

- For sprayer with spray head or trigger separate from spray bottle, first connect

trigger to spray bottle.

- Grasp container and point sprayer away from body.
- Adjust spray nozzle to desired setting.
- Squeeze trigger and direct spray to target weed(s).
- Apply enough product to cover the target weed(s). Spray to wet the weed leaf surface.
- One application should be sufficient.
- Effects begin to show after 24 to 48 hours.
- A repeat application may be required in 3 to 4 weeks for hard to kill (or control) weeds.
- To obtain wider coverage, raise container.

Method

[Spray](#)

Rates

[field_rates 0](#)

[field_rates 1](#)

•

Timings

[Postemergence \(Weed\)](#)

[Apply this product to young, actively growing weeds.](#)