

BEANS (SNAP)

General Information

DIRECTIONS FOR USE:

General Precautions and Restrictions:

DO NOT allow people or pets to enter treated areas until sprays have dried or dusts have settled. DO NOT apply this product in a way that will contact other persons, or pets, either directly or through drift. Only protected handlers may be in the area during application. DO NOT apply this product through any type of irrigation system.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use in small vegetable or fruit gardens, on plants intended for aesthetic purposes or climatic modification and being grown in exterior landscapes, ornamental gardens or parks.

Limitations, Restrictions, and Exceptions

BEANS (SNAP)

Application Directions: Repeat at 7 day intervals.

GARDEN VEGETABLE DISEASES:

Apply FRUIT TREE, VEGETABLE & ORNAMENTAL FUNGICIDE at the rate indicated for each garden crop for the disease to be controlled, by mixing in one gallon of water to cover approximately 200 square feet (10 ft. by 20 ft.) of planted crop. Repeat applications at intervals specified. Use the 7 day schedule during severe disease conditions or wet weather. Do not apply closer to harvest than the indicated Preharvest Interval (PHI) for each type of crop.

Do not apply more than the specified amount of FRUIT TREE, VEGETABLE & ORNAMENTAL FUNGICIDE per 200 sq. ft. of garden area per year for each crop type.

NOTE: Avoid spraying plants during extremely hot and sunny weather. Do not apply to wilted plants. Do not graze treated areas or feed treated plant parts to livestock.

tsp. = teaspoonful (1/6 fluid ounce)

Method

[Spray](#)

Pre-Harvest Interval

7 days

Rates

[field_rates 0](#)

[field_rates 1](#)

•

Timings

[During early bloom stage or when disease first threatens.](#)