

# **BROADLEAF SHRUBS AND TREES, FLOWERING PLANTS AND BULBS, FOLIAGE PLANTS - (LEAFSPOTS AND FOLIAR BLIGHTS)**

General Information

## **DIRECTIONS FOR USE**

### General Precautions and Restrictions

DO NOT allow people or pets to enter treated areas until sprays have dried or dusts have settled. DO NOT apply this product in a way that will contact other persons, or pets, either directly or through drift. Only protected handlers may be in the area during application. DO NOT apply this product through any type of irrigation system.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use in small vegetable or fruit gardens, on plants intended for aesthetic purposes or climatic modification and being grown in exterior landscapes, ornamental gardens or parks.

### Limitations, Restrictions, and Exceptions

#### **BROADLEAF SHRUBS AND TREES, FLOWERING PLANTS AND BULBS, FOLIAGE PLANTS - (LEAFSPOTS AND FOLIAR BLIGHTS)**

**BROADLEAF SHRUBS AND TREES:** (Aspen, Flowering peach, Flowering plum, Magonolia, Maple, Sand Cherry, Sequoia, Spirea and Walnut) Not approved for use in California.

**FLOWERING PLANTS AND BULBS:** (Crocus, Daffodil, Marigold, Narcissus, Pansy, Phlox, Poinsettia, Tulip) Not approved for use in California.

Avoid applications during bloom period on plants where flower injury is unacceptable.

Poinsettia: Discontinue applications prior to bract formation; phytotoxicity is possible on the bracts.

Rose: Use 1½ Tbs. of FRUIT TREE, VEGETABLE & ORNAMENTAL FUNGICIDE per 4 gallons of water.

FOLIAGE PLANTS: (Aglaonema, Areca palm, Artemesia, Boston Fern, Dumbcane, Fatsia, Ficus, Florida ruffle fern, Lipstick plant, Ming Aralia, Peperomia, Zebra plant) Not approved for use in California.

Pachysandra: Use 4 Tbs. of FRUIT TREE, VEGETABLE & ORNAMENTAL FUNGICIDE per 4 gallons of water.

LEAFSPOTS AND FOLIAR BLIGHTS: (Drechslera (Bipolaris) leafspots, inkspot, Gloeosporium black leafspot, Myrothecium leafspot, brown rot, Phyllosticta leafspot, Ramularia leafspot, Stagonospora leaf scorch) Not approved for use in California.

#### ORNAMENTAL GARDEN PLANTS

Apply enough diluted spray to provide thorough coverage of all plant parts that are intended to be protected from disease, generally ranging from ½ to 3 gallons per 1,000 square feet of garden area. Repeat applications at 7 to 10 day intervals until conditions are no longer favorable for disease. During periods when conditions favor severe disease incidence, generally cloudy or wet weather, apply FRUIT TREE, VEGETABLE & ORNAMENTAL FUNGICIDE at 7 day intervals. Fruits and other structures which may be borne on treated plants **MUST NOT BE EATEN**.

Method

[Spray](#)

Rates

[field\\_rates 0](#)

•

Timings

[N.A.](#)