

BROADLEAF SHRUBS AND TREES, FLOWERING PLANTS AND BULBS, FOLIAGE PLANTS - (CYLINDROCLADIUM STEM CANKER)

General Information

DIRECTIONS FOR USE

General Precautions and Restrictions

DO NOT allow people or pets to enter treated areas until sprays have dried or dusts have settled. DO NOT apply this product in a way that will contact other persons, or pets, either directly or through drift. Only protected handlers may be in the area during application. DO NOT apply this product through any type of irrigation system.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use in small vegetable or fruit gardens, on plants intended for aesthetic purposes or climatic modification and being grown in exterior landscapes, ornamental gardens or parks.

Limitations, Restrictions, and Exceptions

BROADLEAF SHRUBS AND TREES, FLOWERING PLANTS AND BULBS, FOLIAGE PLANTS - (CYLINDROCLADIUM STEM CANKER)

BROADLEAF SHRUBS AND TREES: (Eucalyptus) Not approved for use in California.

Cylindrocladium stem canker: Not approved for use in California

ORNAMENTAL GARDEN PLANTS

Apply enough diluted spray to provide thorough coverage of all plant parts that are intended to be protected from disease, generally ranging from ½ to 3 gallons per 1,000 square feet of garden area. Repeat applications at 7 to 10 day intervals until conditions are no longer favorable for disease. During periods when conditions favor severe disease incidence, generally cloudy or wet weather, apply **FRUIT TREE, VEGETABLE & ORNAMENTAL FUNGICIDE** at 7 day intervals. Fruits and other structures which may be borne on treated plants **MUST NOT BE EATEN.**

Method

[Spray](#)

Rates

[field_rates 0](#)

•

Timings

[N.A.](#)