

CONIFER - (AUTOECIOUS NEEDLE RUST (WEIR.S CUSHION RUST) (SPRUCES))

General Information

DIRECTIONS FOR USE

General Precautions and Restrictions

DO NOT allow people or pets to enter treated areas until sprays have dried or dusts have settled. DO NOT apply this product in a way that will contact other persons, or pets, either directly or through drift. Only protected handlers may be in the area during application. DO NOT apply this product through any type of irrigation system.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use in small vegetable or fruit gardens, on plants intended for aesthetic purposes or climatic modification and being grown in exterior landscapes, ornamental gardens or parks.

Limitations, Restrictions, and Exceptions

CONIFER - (AUTOECIOUS NEEDLE RUST (WEIR.S CUSHION RUST) (SPRUCES))

Application Directions: and repeat twice thereafter at 7 to 10 day intervals.

CONIFER DISEASES

Apply FRUIT TREE, VEGETABLE & ORNAMENTAL FUNGICIDE in sufficient water to obtain uniform coverage of tree canopy, including the undersides of leaves, to the point of spray runoff. Mix the indicated rate of FRUIT TREE, VEGETABLE & ORNAMENTAL FUNGICIDE into each 4 gallons of water to be used for each application, generally sufficient to treat 200 square feet. NOTE: DO NOT apply FRUIT TREE, VEGETABLE & ORNAMENTAL FUNGICIDE within one week before or after application of oil or an oil-based pesticide. Do not graze treated areas or feed treated plant parts to livestock.

tsp. = teaspoonful (1/6 fluid ounce)

Method

[Spray](#)

Rates

[field_rates 0](#)

•

Timings

[When 10% of buds have broken.](#)