

# STONE FRUITS

## General Information

### GENERAL INFORMATION

VAPOR GARD is a water emulsifiable organic concentrate for use on plants to reduce water transpiration. The soft, flexible film formed after the spray application dries, will significantly reduce moisture lost by the plant foliage.

All anti-transpirant spray applications must be applied for full coverage. Spray VAPOR GARD for full coverage or use as a dip. No additional spreader is needed with VAPOR GARD. Apply VAPOR GARD at least one hour, during daylight, before an anticipated rain. Sunlight for this time period, is needed for the protective film to set. VAPOR GARD dries on plants to form a clear, glossy film which retards normal moisture loss without interfering with plant growth or normal respiration. VAPOR GARD beautifies plants by polishing leaf surfaces.

### DIRECTIONS

Can Be Used On All Growing Plants Including: Trees (Deciduous and Conifers); Evergreens; Christmas Trees; Shrubs; Turf; Roses; Flowering Plants; Vegetable, Tobacco and Fruit and Nut Trees. Apply VAPOR GARD alone. CAUTION: Do Not apply in spray tank combination with any pesticide, on any crop.

### Limitations, Restrictions, and Exceptions

STONE FRUITS (Such as Peaches, Nectarines and Apricots): To improve color, size and flavor, apply VAPOR GARD at the rate of 1 gallon per acre, 2 to 3 weeks before harvest. The application should be applied in enough water for good coverage, about 200 gallons per acre. The application may hasten maturity by approximately 1 to 3 days. Do Not use in spray tank combination with any pesticide.

### Method

[Spray](#)

Rates

[field rates 0](#)

-

Timings

[Preharvest](#)