

APPLICATION TO ORNAMENTALS

Limitations, Restrictions, and Exceptions

APPLICATION TO ORNAMENTALS

MIDASH 2SC T&O can be applied to ornamental plants in commercial and residential landscapes and interior plantscapes. MIDASH 2SC T&O is a systemic insecticide that is transported within the plant system from the roots to upper foliage. MIDASH 2SC T&O must be applied into a growing area of the plant that allows absorption of the active ingredient. Adding soluble nitrogen type fertilizers to the spray solution when appropriate can promote the uptake of the active ingredient. Application can be made by foliar application or soil applications; including soil injection, drenches, and broadcast sprays. Foliar applications offer locally systemic activity against insect pests. The systemic translocation of active ingredient will be slower when applied to woody plants with soil applications. This delay can take 60 days or longer depending on species and size of plant. To offset this, make applications before anticipated pest infestation. For outdoor ornamentals, broadcast applications cannot exceed a total of 1.6 pints (0.4 lb. of active ingredient) per acre per year.

Method

[Foliar spray](#)

[Drench](#)

[Injection](#)

[Soil application](#)

Rates

[field rates 0](#)

[field rates 1](#)

-

Restricted Entry Interval

12 hours

Exception: If the product is applied by drenching, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

Timings

N. A.