

## **USE SITE: POME FRUITS (ONLY IN AND AROUND RESIDENTIAL AREAS)**

### Limitations, Restrictions, and Exceptions

#### Notes and Restrictions for residential Pome Fruits:

- Apply specified dosage as foliar spray as needed after petal fall is complete.
- For control of rosy apple aphid, apply prior to leafrolling caused by the pest.
- For first generation leafminer control, make first application as soon as petal fall is complete. Greatest leafminer control will result from the earliest possible application. For second and succeeding generations of leafminer, optimal control is obtained from applications made early in the adult flight against egg and early instar larvae. A second application may be required 10 days later if severe pressure continues or if generations are overlapping. A single application may result in suppression only. MIDASH 2SC T&O will not control late stage larvae.
- For San Jose Scale, time applications to the crawler stage. Treat each generation.
- For late season (preharvest) control of leafhopper species, apply MIDASH 2SC T&O while most leafhoppers are in the nymphal stage.
- For optimal control of mealybug, insure good spray coverage of the trunk and scaffolding limbs or other resting sites of the mealybug.
- Do not apply more than 6.0 fluid ounces per acre in a single application. Do not make more than 5 applications.
- Allow 10 or more days between applications. Allow at least 7 days between last application and harvest.
- Not for use in California for control on pears.

#### Method

[Foliar spray](#)

#### Rates

[field\\_rates 0](#)

[field\\_rates 1](#)

[field\\_rates 2](#)

- 

Restricted Entry Interval

12 hours

Exception: If the product is applied by drenching, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

Timings

[After petal fall](#)