

GOLF COURSE TURF DISEASE CONTROL - BERMUDAGRASS DECLINE

General Information

DIRECTIONS FOR USE

Offset 3.6F Fungicide is a soluble concentrate which will control certain pests on the crops listed on the label when applied according to the Directions for Use.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Limitations, Restrictions, and Exceptions

DISEASE CONTROL IN GOLF COURSE TURF

For use on all Golf turf applications of cool season and warm season grasses (such as Bentgrasses, Bluegrasses, Fescues, Ryegrasses, St. Augustine grasses, and Zoysia) or their mixtures. Offset 3.6F Fungicide is not phytotoxic to any of the above mentioned grasses when used in accordance with the label.

Note: Bermudagrass can be sensitive to Offset 3.6F Fungicide under certain conditions. Do not apply consecutive applications during or just after dormancy break. Avoid applications when temperatures are expected to exceed 85 degrees F.

Offset 3.6F Fungicide can be used for the prevention and control of the diseases mentioned in table below. Begin applications when conditions favor disease development and repeat applications as long as these conditions persist.

Preventative treatments can be applied using 28 day intervals as indicated. When treating golf course greens, always treat aprons and approaches. Spray uniformly over the area to be treated with properly calibrated equipment.

Apply the specified amount of Offset 3.6F Fungicide in sufficient water for thorough coverage. A volume of 66 - 132 gallons per acre (1.5 - 3.0 gallons per 1,000 sq ft) is recommended. Apply using properly calibrated low volume, hand held, mechanical

or motorized ground broadcast equipment. Application to small areas may be made with low-pressure handwand or backpack equipment. Maintain constant agitation during application.

Depending on the disease, Offset 3.6F Fungicide should be watered into the crown and active root zone for best results. Make all applications after mowing and allow foliage to dry thoroughly before irrigation. For best results use spray mixture the same day it is prepared.

RESTRICTIONS:

For use on golf course turf only.

- Do not apply this product through any type of irrigation system.
- Do not use on home lawns and turf sites associated with apartment buildings, daycare centers, playgrounds, playfields, recreational park athletic fields, athletic fields located on or next to schools (i.e. elementary, middle and high school), campgrounds, churches, and theme parks.
- Not for homeowner use.
- Not for use on turf being grown for sale or commercial use as sod. Do not use clippings for animal feed.
- Do not exceed 3.6 fl. oz of Offset 3.6F Fungicide per 1,000 sq ft per year.
- Do not apply more than 6 applications per year, except New York.

Notes

Immediately after fungicide is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and root zone of the turf. The amount of water is dependent on the depth of the root zone.

For prevention, begin applications two or four weeks prior to the historical appearance of disease symptoms. Initiate cultural control practices at the same time the fungicide is applied. Refer to your local County Extension Service for this information.

Apply subsequent applications at 28 day intervals.

APPLICATION:

Apply the specified amount of Offset 3.6F Fungicide in 1.5 to 3.0 gallons of water per 1000 sq. ft. Make all applications after mowing and allow foliage to dry thoroughly before irrigation.

RESTRICTIONS:

Do not use clippings for animal feed. Do not exceed 3.6 fl. oz of Offset 3.6F Fungicide per 1000 sq. ft. per year. Do not exceed 6 applications per year. In New York State, do not exceed 3 applications per year.

Method

[Broadcast/Foliar Ground](#)

Rates

[field_rates 0](#)

-

Timings

[When conditions favor disease development.](#)

[Two or four weeks prior to the historical appearance of disease symptoms.](#)