

AZALEAS, CAMELLIAS, RHODODENDRONS AND OTHER LANDSCAPE (ORNAMENTAL) SHRUBS

General Information

DIRECTIONS FOR USE

Offset 3.6F Fungicide is a soluble concentrate which will control certain pests on the crops listed on the label when applied according to the Directions for Use.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Limitations, Restrictions, and Exceptions

DISEASE CONTROL IN CONTAINER ORNAMENTALS AND COMMERCIAL AND RESIDENTIAL LANDSCAPES

Offset 3.6F Fungicide can be used in a preventative and curative disease control program for the listed plant types and disease in the table below. Optimum disease management is obtained when Offset 3.6F Fungicide is used in conjunction with sound disease management practices.

Apply material with properly calibrated hand held, mechanical or motorized spray equipment. Begin applications when disease first appears and repeat at 14-21 day intervals during the growing season. Use the shortest interval when conditions are unusually favorable for the development of disease. For hand held, mechanical, or motorized applications, mix as directed below and apply as a full coverage spray to drip for the prevention and control of the diseases listed below. Choose a finished spray volume appropriate for the size of the plants and amount of foliage, which will provide thorough coverage throughout the canopy. Allow sprays to dry before overhead irrigation is applied.

Apply Offset 3.6F Fungicide at rates of 4-10 fl oz per acre in 100 gallons of water. Spray volume may range from 50 up to 300 gallons of finished spray per acre depending upon equipment, plant species and plant growth stage at time of

application.

Note: The “Directions for Use” of this product reflect the cumulative inputs from both historical field use and product testing programs. However, it is impossible to test this product on all species and cultivars. A preliminary trial is suggested on a small scale before a full treatment is applied to any plant type not shown on the label but found in a similar use site with a listed disease problem. Wait 5-7 days after treatment to evaluate results. This product is not recommended for use on African Violets, Begonias, Boston Fern, and Geraniums.

ORNAMENTAL USE RESTRICTIONS AND PRECAUTIONS

For use on ornamental plants only; not for woodlands or forest management.

- Not for homeowner use.
- Do not apply more than 10 fl oz per acre in a single application.
- Do not apply more than 0.94 gallons (120 fl oz) of Offset 3.6F Fungicide (equal to 3.38 lbs of tebuconazole) per acre per year.
- Do not make more than 4 applications per year at highest rate.
- Do not apply to bearing fruit trees or vegetables.

TO PREVENT DISEASES - Petal Blight—Apply 2-3 times per week into the flowers as they open and develop color.

TO TREAT EXISTING DISEASE - Apply every 14 days for a total of 3 applications beginning at the first sign of disease.

HOW MUCH TO USE FOR SMALL PLANTINGS: ADD 1 TEASPOON TO 2.5 GALLONS OF WATER.

Pump Style Sprayers

1. Add the appropriate amounts of concentrate and water to the sprayer tank.
2. Close the sprayer, shake well and pressurize.
3. Adjust nozzle to a coarse spray pattern and apply.

4. Occasionally re-pressurize the sprayer if needed to maintain a good spray pattern.

Method

[Spray](#)

Rates

[field rates 0](#)

•

Timings

[At the first sign of disease.](#)