

LANDSCAPE ORNAMENTALS

General Information

GENERAL INFORMATION

This product is a herbicide that provides control of crabgrass and other annual grasses and broadleaf weeds in established lawns and ornamental turfs including golf course fairways, roughs and tee boxes. This product also provides post-emergent crabgrass control up to the 4-leaf stage. This product will not harm nearby established ornamentals when used according to label directions.

Limitations, Restrictions, and Exceptions

LANDSCAPE ORNAMENTALS

16-8-8 with 0.20% Dimension provides preemergence control of listed annual grass and broadleaf weeds in landscape ornamentals. It may be applied to soil surfaces or mulch where ornamental plants are grown for aesthetic purposes in ornamental gardens, parks, golf courses and residential properties. Ornamental plants on the label have shown tolerance to 16-8-8 with 0.20% Dimension when applied as directed. It is impossible, however, to evaluate tolerance for all ornamental plant species or varieties or under all possible growing conditions. Under your growing conditions, a limited area involving only a few plants should be treated for tolerance evaluation prior to large-scale applications.

USE RATES

Apply 16-8-8 with 0.20% Dimension with a properly calibrated spreader that will assure accurate, even particle distribution. This product may be applied as a single application or as split application. Refer to the following table for product rates when treating a small area.

DO NOT apply more than 5.74 lbs. of this product/1,000 sq. ft. per application, and no more than 13.7 lbs. of this product per 1,000 sq. ft. per year.

In New York State, this product may only be used by Professional Applicators at no more than 250 lbs. per acre (5.74 lbs./1,000 sq. ft.), (0.5 lbs. active ingredient per

acre) per year; and is prohibited from use in Nassau and Suffolk Counties.

APPLICATION TIMING

16-8-8 with 0.20% Dimension is a preemergence herbicide that will not control established weeds. Make applications prior to weed seed germination or mulch or to bare ground. The best weed control is obtained when applied to soil that is free of clods, weeds and debris such as leaves. Prior to making an application, control existing vegetation by hand weeding, cultivation or with a postemergence herbicide. Once an application of this product has been made, do not disturb the soil surface as the herbicide barrier will be broken.

PRECAUTIONS

- Only apply this product to established ornamentals.
- DO NOT apply this product directly to bare root of ornamental plants as injury may result.
- DO NOT incorporate this product into soil. Dilution of active ingredient and possible injury to plant roots may occur.
- DO NOT apply to soil around ornamental plants that are under stress due to drought, flooding, excessive fertilizer or soil salts, wind injury, hail, frost damage, winter injury, injury from previously applied pesticides or injury due to insects, nematodes or disease.
- To avoid foliar damage to ornamentals, irrigate immediately after application to remove herbicide particles from the foliage.
- DO NOT apply under conditions that would affect product distribution. Wind could affect distribution and move product onto sensitive species. Uneven product distribution will result in uneven weed control.
- DO NOT apply this product on grasses grown for seed.
- DO NOT graze livestock or feed forage cut from areas treated with this product.

TOLERANT ORNAMENTALS

When applied as directed under the conditions described on the label, the

ornamentals listed below have shown tolerance in field trials. However, this product has not been tested on all cultivars of each species or under all possible growing conditions. Follow directions given above to determine plant tolerance under your growing conditions prior to large-scale use.

Ornamental species only: DO NOT USE ON FOOD PRODUCING TREES AND ORNAMENTALS.

APPLICATION EQUIPMENT AND INSTRUCTIONS

Apply this product with drop or rotary-type spreaders designed to apply granular herbicides. Avoid the use of spreaders that would apply this product in narrow rows or concentrated bands. Before each application, calibrate the spreader according to the equipment manufacturer's directions for adjusting the spreader settings such that the spreader delivers the appropriate application rate recommended above. Apply this product uniformly over the area to be treated. A more uniform application can be made by spreading half the required amount over the area and applying the remaining half at a right angle to the previous direction. Avoid streaking, skips, or overlaps during application. Check equipment frequently to ensure equipment is functioning properly and delivering uniform distribution of granules.

Method

[Broadcast](#)

Rates

[field rates 0](#)

[field rates 1](#)

[field rates 2](#)

•

Timings

[Preemergence \(Weed\)](#)