

# **POTATOES**

## General Information

### GENERAL USE DIRECTIONS

**DILUTION RATE:** Unless otherwise specified, use the rate per acre and dilute as follows:

Air: 5-30 gallons of water per acre.

Dilute: 100 plus gallons of water per acre.

Concentrate: 10-100 gallons of water per acre.

Do not exceed 2 lbs. of product per gallon of water.

Add approximately 1/2 water to tank before adding Folo Spray 20-20-20, agitate thoroughly while adding remaining water, add pesticide last.

When temperatures are cold allow extra time for the product to completely dissolve. Do not use when temperatures are expected to exceed 90°F or when plants are under stress conditions such as heat or water stress.

**THE PRODUCT WILL LOWER THE pH OF THE SPRAY SOLUTION.**

**CAUTION:** The product contains Boron and Molybdenum. Use of the product at rates other than those recommended, may cause injury to susceptible plants.

**DO NOT EXCEED RECOMMENDED AMOUNTS PER ACRE.**

## Limitations, Restrictions, and Exceptions

### POTATOES

- For best results, apply when plants are young and continue at 7 to 10 day intervals through 30 day tuber setting period.

## Method

### [Spray](#)

Rates

[field\\_rates 0](#)

- 

Timings

[At initial tuber formation.](#)