

SOIL APPLICATION - ORCHARD AND VINEYARDS

General Information

DIRECTIONS FOR USE

Actagro Micro Booster can be applied to most vegetable and cucurbit crops, field and row crops, deciduous fruit and nut trees, grapes, berries, olives, avocados and citrus. Micro Booster is designed for soil application and for foliar use. Apply directly to soils or inject through irrigation systems where water quality is satisfactory to maintain solubility. Foliar applications should be as a dilute spray. Application rates vary by crop or situation. Mixing with irrigation water for field storage is generally not recommended. Products blended with irrigation water should be used immediately and not stored for prolonged periods.

GENERAL INFORMATION

Micro Booster will be influenced by various soil and environmental conditions and should be applied under conditions of optimum plant growth. Optimum conditions are defined as adequate humidity, moderate temperatures and adequate soil moisture. Optimum conditions for most crops generally occur during early morning or late evening hours.

CAUTION: If any other chemicals are to be used in the spray mixture be sure to check compatibility. Foliar application of Micro Booster close to harvest may leave a brown residue on the harvestable portion of the crop. Rainfall, sprinkler irrigation or high humidity should alleviate this condition.

Limitations, Restrictions, and Exceptions

SOIL APPLICATION

Micro Booster may be applied pre-plant, as a starter, side dress, fertigated, top dressed or banded to the soil. To apply alone, dilute with water to ensure uniform distribution. Repeat applications may be beneficial.

ORCHARDS & VINEYARDS: Inject 1 to 4 gallons per acre 3 to 5 inches deep at the drip line, or surface band and water into the soil. Repeat applications may be beneficial.

Method

[Band](#)

[Injection](#)

[Soil application](#)

Rates

[field_rates 0](#)

•

Timings

[N.A.](#)