

SOIL APPLICATION - ALL CROPS

General Information

GENERAL INFORMATION

RESIST II contains potassium, which is intended to stimulate root development and resistance to disease.

RESIST II is non-phytotoxic when used as directed and is compatible with most commonly used pesticides and fertilizers.

RESIST II has a low salt index and can be applied as a foliar or used in drip or micro-sprinkler irrigation systems.

RESIST II should not be applied to plants that are stressed by high temperature or lack of moisture. Phosphorous acid products are intended to be used as a supplemental fertilizer treatment.

* When using RESIST II as a foliar application, the phosphite ions are taken up directly by the plant foliage and may undergo a degree of conversion to phosphate ions, or are utilized directly by plants, as phosphite ions. When using RESIST II as a soil application on annual crops, you may experience a lesser response from the initial crop, with a corresponding superior response from succeeding crops. In addition, placement close to seeds or root zones may be injurious to crops. The effect may be aggravated by a soil pH below 6.5.

DIRECTIONS FOR USE

RESIST II can be applied to most vegetable and cucurbit crops, field crops, row crops, citrus, fruit and nut trees, vines, olives, turf, nursery and greenhouse plants, and ornamental farms. RESIST II may be applied to the soil or used as a foliar and may be applied through center pivot irrigation systems.

Limitations, Restrictions, and Exceptions

ALL CROPS: Apply to the soil at 1 quart to 2 quarts per acre. RESIST II is most effective with drip or micro-sprinkler irrigation systems. Repeat as needed.

Method

[Soil application](#)

Rates

[field rates 0](#)

-

Timings

[N.A.](#)