

FOLIAR APPLICATION - ALL CROPS

General Information

GENERAL INFORMATION

Resist SW is non-phytotoxic when used as directed and is compatible with most commonly used pesticides and fertilizers. Resist SW has a low salt index and can be applied as a foliar or used in drip or micro-sprinkler irrigation systems. Resist SW should not be applied to plants that are stressed by high temperature or lack of moisture.

CAUTION: Always check for compatibility with other products prior to use. Do not mix with conventional calcium based fertilizers. Actagro Resist SW is not compatible with conventional products containing the salts of heavy metals.

CAUTION: Foliar application of Resist SW close to harvest may leave a brown residue on the harvestable portion of the crop. Rainfall, sprinkler irrigation or high humidity should alleviate this condition.

DIRECTIONS FOR USE

Resist SW can be applied to most vegetable and cucurbit crops, field and row crops, deciduous fruit and nut trees, grapes, berries, olives, avocados and citrus. Resist SW is designed for soil application and for foliar use. Apply directly to soils or inject through irrigation systems where water quality is satisfactory to maintain solubility. Foliar applications should be as a dilute spray. Application rates vary by crop or situation. Mixing with irrigation water for field storage is generally not recommended. Products blended with irrigation water should be used immediately and not stored for prolonged periods.

Limitations, Restrictions, and Exceptions

FOLIAR APPLICATION

ALL CROPS: Resist SW may be applied by air or ground spray. Use the appropriate volume of water for the application method and the crop. Use a minimum of 10 gallons of water per acre. Additional applications may be necessary.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field_rates 0](#)

-

Timings

[N.A.](#)