

# **APPLES AND MAYHAWES - SCAB - POSTINFECTION**

## General Information

### USE INFORMATION

Read all Directions for Use carefully before applying.

SONOMA 40WSP fungicide is a systemic, protectant and curative fungicide recommended for the control of specific diseases mentioned on the label. Optimum disease control is achieved when the fungicide is applied in a regularly scheduled preventative spray program.

### Use Rate Determination

Carefully read, understand and follow label use rates and restrictions. Under low disease conditions, minimum label use rates per application can be used while maximum label rates and shortened spray schedules are recommended for severe or threatening disease conditions. For proper application, determine the number of acres to be treated, the recommended label use rate and the gallonage to be applied per acre. Prepare only the amount of spray solution required to treat the measured acreage. Careful calibration of spray equipment is recommended prior to use.

### USE DIRECTIONS FOR TREE FRUITS

Best control of labeled diseases is achieved when SONOMA 40WSP is applied on a 7- to 10-day application schedule.

SONOMA 40WSP is a systemic fungicide and does not redistribute with rainfall after application. Application equipment spray nozzles should be adjusted to apply a uniform spray throughout the entire tree canopy. The following use recommendations are to be used as guidance in determining the amount of SONOMA 40WSP to be used per 100 gallons spray or per acre. Refer to specific tree fruit use directions to determine actual use rates for the control of labeled diseases.

Dilute, Thorough Coverage Applications: Dilute thorough coverage applications are based on the amount of spray solution required to thoroughly wet trees until spray runoff. The following specific use directions for apples and mayhaws utilizes a 400

gallon per acre dilute basis and the specific use directions for stone fruits utilize a 250 gallon per acre dilute basis. Thorough spray coverage is essential for good disease control. SONOMA 40WSP should be applied in sufficient water to ensure adequate coverage.

Limitations, Restrictions, and Exceptions

## APPLICATION DIRECTIONS

SONOMA 40WSP provides 96-hour post-infection control or curative activity. Apply as soon as possible after infection period. Follow with a standard preventative spray schedule.

- Based on dilute sprays with a 400 gallon per acre basis.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

14 days

Rates

[field rates 0](#)

[field rates 1](#)

•

Restricted Entry Interval

24 hours

Timings

[Postinfection](#)