

## **APPLICATION TO CONSERVATION RESERVE PROGRAM (CRP) ACRES**

### General Information

#### PRODUCT INFORMATION

ALLIGARE FLUROXYPYR HERBICIDE is a selective herbicide for control of annual and perennial broadleaf weeds including kochia, bedstraw (cleavers), common sunflower, volunteer flax and volunteer potato on-farm non-cropland, non-cropland areas including industrial sites, nonirrigation ditch banks, rights of way, pine plantations and conservation reserve program (CRP).

#### SPOT TREATMENTS AND HAND-HELD SPRAYERS

To prevent misapplication, spot treatments should be applied with a calibrated boom or with hand sprayers according to the directions below.

Hand-held or backpack sprayers may be used for spot application of ALLIGARE FLUROXYPYR HERBICIDE if care is taken to apply the spray uniformly and at a rate equivalent to a broadcast application. Application rates in the table are based on an area of 1,000 sq. ft. The amount of ALLIGARE FLUROXYPYR HERBICIDE (fl oz or ml) in the table should be mixed with 1 gallon or more of water and applied to an area of 1,000 sq ft. To calculate the amount of product required for larger areas, multiply the table value (fl oz or ml) by the area to be treated in “thousands” of square feet, e.g. if the area to be treated is 5,000 sq ft, multiple the table value by 5 (calc.  $5,000 / 1000 = 5$ ).

#### Limitations, Restrictions, and Exceptions

#### APPLICATION TO CONSERVATION RESERVE PROGRAM (CRP) ACRES

Do not use on CRP acres that are underseeded with desirable legumes, clovers, or other sensitive broadleaf plants. ALLIGARE FLUROXYPYR HERBICIDE may be applied to CRP acres. For best results, apply as a single broadcast post emergence treatment using ground equipment or by air to control susceptible broadleaf weeds. Apply at a rate of 0.4 to 0.7 pt/A when weeds are small and actively growing, but before weeds are 8 inches tall or vining. Spot treatments should be applied at rates

and spray volumes equivalent to broadcast application. See instructions for “Spot Application” above.

Restriction: Grazing or haying of treated CRP acres is prohibited.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Spot treatment](#)

Rates

[field\\_rates 0](#)

- 

Restricted Entry Interval

24 hours

Timings

[Postemergence \(Weed\)](#)