

## **GRAPES AND BERRIES**

### General Information

#### GENERAL INFORMATION

Metalosate Calcium is designed for foliar application on plants to prevent or correct nutrient deficiencies that may limit crop growth and yields. It is water soluble and nontoxic to plants when applied as directed. For best results, apply Metalosate Calcium according to recommendations based on plant tissue or soil analysis.

#### DIRECTIONS:

Dilute one part Metalosate Calcium with 20 or more parts water. Apply by a spray method and in an adequate amount of water that will provide complete coverage of the plants. Use of a non ionic wetting agent may improve spray coverage of certain hard to wet plants. Do not apply undiluted.

Metalosate Calcium can be included in a regular spray program on crops. Consult with an Albion representative on compatibility with other spray materials.

The rate of application will depend on the crop, stage of growth, and severity of deficiency. The maximum recommended rates are for mature, full sized plants. Reduce the rates proportionately when spraying smaller plants.

Metalosate Calcium may be used on legumes, grain crops, root crops, cucurbits, cole crops, leafy vegetables, woody and herbaceous ornamentals, deciduous fruits, vine crops, tropical and subtropical fruits, and many other crops.

#### Limitations, Restrictions, and Exceptions

#### GRAPES AND BERRIES

Make an application of 16 to 32 fluid ounces per acre after active growth begins. The application may be repeated at intervals of one week or more through the vegetative growth period.

#### Method

[Foliar spray](#)

Rates

[field\\_rates 0](#)

- 

Timings

[After active growth begins.](#)