

FIELD CORN, SWEET CORN, ETC. - IN-FURROW

General Information

SmartBox Information: AZTEC 4.67 G must be applied with the SmartBox system. Refer to the SmartBox equipment and use manual for information on system operation and calibration.

AZTEC 4.67 G is not systemic and will not interact with corn herbicides. It will not affect seed germination or crop emergence when applied in-furrow.

OPTIMUM CONTROL - Optimum control of cutworm is achieved when AZTEC 4.67 G is applied as a Band or T-Band.

Optimum control of wireworms, seedcorn maggot, seedcorn beetle and white grubs is achieved when AZTEC 4.67 G is applied as a T-Band or In-Furrow.

For no-till corn where runoff drains into adjacent or neighboring aquatic sites (lakes, reservoirs, rivers, permanent streams, marshes, estuaries, natural ponds, or ponds used for commercial fishing), apply AZTEC 4.67 G as an in-furrow or T-band application only.

For applications made within 20 yards of aquatic sites (lakes, reservoirs, rivers, permanent streams, marshes, estuaries, natural ponds, or ponds used for commercial fishing), apply as an in-furrow application only.

Cover or incorporate spills (including end row spillage).

IMPORTANT: Read the entire directions for use, maintenance sections for the SmartBox System and these entire.

Directions and Conditions of Sale for AZTEC 4.67 G prior to use.

For use only in the states of: Arkansas, Colorado, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Michigan, Minnesota, Missouri, Montana, Nebraska, Nevada, New Mexico, North Dakota, Ohio, Oklahoma, Pennsylvania, South Dakota, Tennessee, Utah, Vermont, West Virginia, Wisconsin, and Wyoming.

Do not use on other crops grown for food or forage. Corn may be replanted anytime after application of AZTEC 4.67 G. All other crops may be planted 30 days after

application.

Limitations, Restrictions, and Exceptions

IN-FURROW – Apply specified dosage at planting directly into the seed furrow behind the planter shoe and in front of the press wheel.

A maximum of 3.27 pounds AZTEC 4.67 G may be applied per acre per crop season.

Row Spacing (inches) - 30-40

Pounds Per Acre - 3.27-2.45

Method

[In Furrow](#)

Rates

[field rates 0](#)

-

Restricted Entry Interval

48 hours

Exception: If the product is soil injected or soil incorporated, the Worker Protection Standard under certain circumstances, allows workers to enter the treated area without restrictions if there will be no contact with anything that has been treated.

Timings

[At-Plant](#)

[Postemergence \(Crop\)](#)

[Preemergence \(Crop\)](#)

[Seed germination.](#)