

FIELD CORN, POPCORN AND SWEET CORN - AT PLANTING

General Information

DIRECTIONS FOR USE

All setbacks must be planted to a crop or seeded with grass or other suitable cover, and all vegetative buffers must be seeded with grass or other suitable cover.

DO NOT load, rinse, or wash equipment within 300 ft from surface water bodies.

DO NOT load, rinse, or wash equipment within 50 ft from wells unless conducted on an impervious surface.

To prevent channeling of surface water runoff, adjust the planter row-cleaners appropriately to prevent rows lower in height than adjacent soil.

Limitations, Restrictions, and Exceptions

FIELD CORN, POPCORN AND SWEET CORN - AT PLANTING

Application

Banded

Place granules in a 7 inch band over the row, in front of the presswheel and incorporate evenly into the top 1 inch of the soil.

In-Furrow

Place granules directly in the seed furrow behind the planter shoe.

Remarks

For use on conventional, conservation, or no-till corn. In situations where crop debris or winds can prevent proper placement of granules, in-furrow applications are recommended. Infurrow applications reduce the potential for insecticide runoff resulting from excessive rain. The presswheel should be adjusted so that the closed seed furrow is higher than the surrounding soil to reduce the potential for insecticide run-off resulting from excessive rain.

DO NOT exceed 8.7 pounds of Counter 15G per acre.

Suppression of: Cutworms, Lesser cornstalk borers.

Corn rootworms - Counter 15G controls species of corn rootworms including Southern, Northern, Western, and Mexican corn rootworms.

Chinchbugs - For use in the states of Alabama, Florida, Georgia, Louisiana, Mississippi and Texas only for early season control of light to moderate infestations.

Cutworms - Under dry soil conditions or heavy infestations, it may be necessary to apply an insecticide rescue treatment with another registered insecticide after corn emergence to control surviving cutworm larvae.

The 6.0 oz. rate should only be used on first-year corn. Use the high rate on first-year corn where heavy rootworm population pressures are anticipated or for control of other pests.

If application is made at planting, do not make postemergence or cultivation time treatments of Counter 15G

ALS-inhibiting herbicides SHOULD NOT be used if Counter 15G has been applied to corn at the time of planting. See additional information for “at cultivation” applications.

ALS-inhibiting herbicides such as Accent, Accent Gold, Basis Gold, Steadfast, Beacon, Exceed, Hornet™, Permit, Scorpion III, Celebrity, Celebrity Plus, Lightning, Northstar, and Spirit herbicides.

Method

[In Furrow](#)

[Band](#)

Rates

[field_rates 0](#)

•

Restricted Entry Interval

48 hours

The 48-hour REI is increased to 72 hours in outdoor areas where average rainfall is less than 25 inches a year.

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

Timings

[At-Plant](#)