

SUGAR BEETS - APHIDS, SUGAR BEET CYST NEMATODE AND LEAFHOPPER (AT PLANTING)

General Information

DIRECTIONS FOR USE

All setbacks must be planted to a crop or seeded with grass or other suitable cover, and all vegetative buffers must be seeded with grass or other suitable cover.

DO NOT load, rinse, or wash equipment within 300 ft from surface water bodies.

DO NOT load, rinse, or wash equipment within 50 ft from wells unless conducted on an impervious surface.

To prevent channeling of surface water runoff, adjust the planter row-cleaners appropriately to prevent rows lower in height than adjacent soil.

Limitations, Restrictions, and Exceptions

SUGAR BEETS - APHIDS, SUGAR BEET CYST NEMATODE AND LEAFHOPPER (AT PLANTING)

Application

Knifed-in

Drill granules 2 inches into the side of the seed and 2 to 4 inches below the seed.

Remarks

At Planting

DO NOT place granules in direct contact with the seed or crop injury may occur. Power incorporation may be used.

DO NOT incorporate deeper than 2 inches.

Only one application of Counter 15G per year may be made.

DO NOT harvest sugar beets or feed tops to livestock within 150 days of a knifed-in

application

For any row spacing (minimum 20 inch row spacing) or no more than 13.1 lbs. of Counter 15G per acre.

Method

[N.A.](#)

Pre-Harvest Interval

150 days

Rates

[field_rates 0](#)

-

Restricted Entry Interval

48 hours

The 48-hour REI is increased to 72 hours in outdoor areas where average rainfall is less than 25 inches a year.

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

Timings

[At-Plant](#)