SUGAR BEETS - APHIDS, SUGAR BEET CYST NEMATODE AND LEAFHOPPER (POSTEMERGENCE)

General Information

DIRECTIONS FOR USE

All setbacks must be planted to a crop or seeded with grass or other suitable cover, and all vegetative buffers must be seeded with grass or other suitable cover.

DO NOT load, rinse, or wash equipment within 300 ft from surface water bodies.

DO NOT load, rinse, or wash equipment within 50 ft from wells unless conducted on an impervious surface.

To prevent channeling of surface water runoff, adjust the planter row-cleaners appropriately to prevent rows lower in height than adjacent soil.

Limitations, Restrictions, and Exceptions

SUGAR BEETS - APHIDS, SUGAR BEET CYST NEMATODE AND LEAFHOPPER (POSTEMERGENCE)

Application

Banded

Apply in 5 to 7 inch band over the row and lightly incorporate into the soil with cultivation shovels or other suitable implement.

Remarks

Postemergence

For sugar beet root maggot control apply at the time of first fly emergence.

DO NOT harvest sugar beets or feed tops to livestock within 110 days after at planting banded, at planting modified in-furrow or postemergence banded applications.

DO NOT harvest sugar beets or feed tops to livestock within 150 days of a knifed-in

application

Crown borer - Postemergence - apply Counter 15G at the first sign of crown borer infestation.

For any row spacing (minimum 20 inch row spacing)

Method

In Furrow

Band

In Furrow

Band

In Furrow

Band

Pre-Harvest Interval

110 days

Rates

field rates 0

•

Restricted Entry Interval

48 hours

The 48-hour REI is increased to 72 hours in outdoor areas where average rainfall is less than 25 inches a year.

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

Timings

At-Plant

Postemergence (Crop)

At the time of first fly emergence.