

GRAIN SORGHUM - AT PLANTING

General Information

DIRECTIONS FOR USE

All setbacks must be planted to a crop or seeded with grass or other suitable cover, and all vegetative buffers must be seeded with grass or other suitable cover.

DO NOT load, rinse, or wash equipment within 300 ft from surface water bodies.

DO NOT load, rinse, or wash equipment within 50 ft from wells unless conducted on an impervious surface.

To prevent channeling of surface water runoff, adjust the planter row-cleaners appropriately to prevent rows lower in height than adjacent soil.

Limitations, Restrictions, and Exceptions

GRAIN SORGHUM - AT PLANTING

Application

Banded

Place granules in a 5 to 7 inch band directly behind the planter shoe in front of the press wheel and lightly incorporate with drag chains or tines.

Banded

DO NOT use banded applications for aphid control in New Mexico, west Texas and the panhandle of Oklahoma.

Remarks

DO NOT graze or harvest grain sorghum forage within 50 days after treatment.

DO NOT harvest grain sorghum fodder or grain within 100 days after treatment

Chinchbugs - For use in the states of Alabama, Arkansas, Florida, Georgia, Louisiana, Mississippi and Texas only for early season control of light to moderate infestations.

For any row spacing (minimum 20 inch row spacing) or no more than 11.3 lbs. of Counter 15G per acre.

Method

[Band](#)

Rates

[field_rates 0](#)

-

Restricted Entry Interval

48 hours

The 48-hour REI is increased to 72 hours in outdoor areas where average rainfall is less than 25 inches a year.

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

Timings

[At-Plant](#)