

# **ASPARAGUS**

## General Information

### GENERAL INFORMATION

### SMARTBOX INFORMATION

The SmartBox system should be properly calibrated by a person familiar with operation of the system to assure an accurate application rate of Lorsban 15G. Lorsban 15G must be applied with the SmartBox system. Please refer to the SmartBox operator's manual for detailed information on system operation and calibration procedures. Do not attempt to open or tamper with the SmartBox container. Do not attempt to repair the SmartBox container. If repairs are necessary, contact your local AMVAC sales representative or the manufacturer of the SmartBox system.

Refer to the label for the Application Rates/1000 Ft of Row and Equivalent/Acre at Different Row Spacings.

### Limitations, Restrictions, and Exceptions

### ASPARAGUS

Postharvest Treatment: Apply Lorsban 15G as a postharvest ground application for suppression of the target pest. Apply as a band over the entire crown area when the asparagus beds have been split (i.e., remove most of the soil from above the asparagus crowns). Cover the area with soil the day of application. Note: Control may be reduced in soils with high organic matter content.

### Specific Use Restrictions:

For use only in the Midwest and Pacific Northwest states.

### Method

[Broadcast/Foliar Ground](#)

[Band](#)

Pre-Harvest Interval

180 days

Rates

[field\\_rates 0](#)

- 

Restricted Entry Interval

24 hours

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

Timings

[Post-harvest](#)