

CITRUS ORCHARD FLOORS

General Information

GENERAL INFORMATION

SMARTBOX INFORMATION

The SmartBox system should be properly calibrated by a person familiar with operation of the system to assure an accurate application rate of Lorsban 15G. Lorsban 15G must be applied with the SmartBox system. Please refer to the SmartBox operator's manual for detailed information on system operation and calibration procedures. Do not attempt to open or tamper with the SmartBox container. Do not attempt to repair the SmartBox container. If repairs are necessary, contact your local AMVAC sales representative or the manufacturer of the SmartBox system.

Refer to the label for the Application Rates/1000 Ft of Row and Equivalent/Acre at Different Row Spacings.

Limitations, Restrictions, and Exceptions

CITRUS ORCHARD FLOORS

Broadcast Treatment: To control foraging ants and suppress mounds, apply Lorsban 15G in combination with a rotary granular applicator, such as a fertilizer spreader, that will uniformly broadcast the granules over the orchard floor. Do not apply where weed growth or other obstructions would impede uniform coverage of the orchard floor. Foliar applications of Lorsban 4E may be made in addition to orchard floor treatments of Lorsban 15G.

Specific Use Restrictions:

- Do not make a second application within 10 days of any application of chlorpyrifos to the orchard.
- Do not allow meat or dairy animals to graze in treated areas.

Method

Broadcast Application

Pre-Harvest Interval

28 days

Rates

field rates 0

-

Restricted Entry Interval

24 hours

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

Timings

Post-harvest