

RADISH - ROOT MAGGOTS

General Information

GENERAL INFORMATION

SMARTBOX INFORMATION

The SmartBox system should be properly calibrated by a person familiar with operation of the system to assure an accurate application rate of Lorsban 15G. Lorsban 15G must be applied with the SmartBox system. Please refer to the SmartBox operator's manual for detailed information on system operation and calibration procedures. Do not attempt to open or tamper with the SmartBox container. Do not attempt to repair the SmartBox container. If repairs are necessary, contact your local AMVAC sales representative or the manufacturer of the SmartBox system.

Refer to the label for the Application Rates/1000 Ft of Row and Equivalent/Acre at Different Row Spacings.

Limitations, Restrictions, and Exceptions

RADISH

At Plant In-Furrow Treatment: Apply the granules in the seed furrow at planting time so that the granules are covered with soil.

Specific Use Restrictions:

The maximum single application rate is 0.5 oz active chlorpyrifos per 1000 feet of row (2.76 lb active chlorpyrifos per acre).

Method

[In Furrow](#)

Pre-Harvest Interval

7 days

Rates

[field_rates 0](#)

-

Restricted Entry Interval

24 hours

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

Timings

[At-Plant](#)