

FOR THE CONTROL OF FERAL PIGEONS

General Information

PREBAITING DIRECTIONS

This product must not be applied where non-target birds are feeding. Careful observations of the birds' feeding habits must therefore be made to establish proper feeding locations and to determine that no non-target birds are feeding on the prebait. Prebaiting with untreated grains of the same composition as the Avitrol carrier is usually essential to the effective deterrence of birds with Avitrol. The exact prebaiting technique employed will vary depending on the birds' location and the surrounding environment. The locale should be studied for a period long enough to ascertain the birds' habits in the area. Proper prebaiting and treatment with diluted Avitrol grain will minimize the chance of bird mortality while producing the optimum conditions for the desired results. Pigeons will feed from roof tops as well as from the ground. Feed acceptance from trays will normally require more time.

DILUTION DIRECTIONS

Avitrol treated grain should be thoroughly mixed with untreated grain of the same composition as the Avitrol carrier at the dilution ratios indicated below. For the control of Pigeons with Avitrol, a dilution ratio of one part treated to twenty-nine parts untreated can be used in most situations. However, where other bird food is available, this ration may need to be adjusted to a one part treated to nineteen or fourteen parts untreated, or even lower. No dilution ratio less than one part treated to nine parts untreated is recommended.

Limitations, Restrictions, and Exceptions

FOR THE CONTROL OF FERAL PIGEONS

BAITING DIRECTIONS

To obtain minimal mortality, the distribution of Avitrol should be limited to scattered spot placements that will provide feeding opportunities only for the necessary number of target birds. After the birds' feeding patterns have been established through prebaiting, replace untreated bait with diluted treated bait only at sites where the target birds are actively feeding sites. Do not apply treated bait to

inactive feeding sites. Where uneaten bait may be a hazard to other birds or animals, it should be picked up at the end of each day. Pick up and dispose of dead birds by burial. This product must not be applied where nontarget bird species are feeding. During the first few days, baiting with Avitrol may require successive applications until control is obtained. Note that retreatment may also necessary after rainfall.

Method

[Bait](#)

Rates

[field_rates 0](#)

•

Timings

[N.A.](#)