

FIELD EQUIPMENT DISINFECTION

General Information

DIRECTIONS FOR USE

SaniDate 5.0 Sanitizer/Disinfectant works best when diluted with water containing minimal levels of organic or inorganic materials, and with water having a neutral pH. Thoroughly rinse out tank with water before mixing concentrate. This product will readily mix with clean, neutral water and does not require agitation.

SaniDate 5.0 Sanitizer/Disinfectant concentrate should not be combined or mixed with any other pesticide concentrates.

This product is not for use on medical device surfaces.

Before use in federally inspected meat and poultry food processing plants and dairies, food products and packaging materials must be removed from room or carefully protected. A potable water rinse is not allowed (Do not rinse) following use of the product as a sanitizer on previously cleaned hard surfaces provided that the surfaces are adequately drained before contact with food so that little or no residue remains.

Limitations, Restrictions, and Exceptions

FIELD EQUIPMENT DISINFECTION

SaniDate 5.0 Sanitizer/Disinfectant may be used to disinfect harvest equipment such as pickers, harvesters, trailers, trucks (including truck body parts and tires), bins, packing crates, ladders, power tools, hand tools, gloves, rubber boots, pruning shears or other equipment that may transfer *Xanthomonas axonopodis* (citrus canker surrogate). This product can also be used to disinfect surfaces contaminated with *Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Salmonella enterica*, *Pediococcus damnosus*, and *Lactobacillus malefermentans*. When applied at recommended disinfectant rates, SaniDate 5.0 Sanitizer/Disinfectant is also effective as a fungicide against *Trichophyton mentagrophytes*.

1. Before disinfection, move the field equipment into an area with an impervious

surface and with controlled drainage. Ensure that no disinfection solution will be released into the environment.

2. Remove gross contamination with a cleaner or other suitable detergent and rinse with water.

3. For *Xanthomonas axonopodis*, use SaniDate 5.0 Sanitizer/Disinfectant at a dilution rate of 1:1,000 or 0.65 fl. oz per 5 gallons of water, as a general coarse spray. For bacteria such as *P. aeruginosa*, *S. enterica*, and *S. aureus*, use SaniDate 5.0 Sanitizer/Disinfectant at a dilution rate of 1:256 (0.5 fl. oz/gal) as a general coarse spray.

4. Allow solution to contact surface for ten (10) minutes.

5. Allow to air dry, do not rinse.

Method

[Spray](#)

Rates

[field_rates 0](#)

•

Timings

[N.A.](#)