

# **FOR DIRECT INJECTION INTO SPRAY WATERS USED ON PROCESS LINES**

## General Information

### DIRECTIONS FOR USE

SaniDate 5.0 Sanitizer/Disinfectant works best when diluted with water containing minimal levels of organic or inorganic materials, and with water having a neutral pH. Thoroughly rinse out tank with water before mixing concentrate. This product will readily mix with clean, neutral water and does not require agitation.

SaniDate 5.0 Sanitizer/Disinfectant concentrate should not be combined or mixed with any other pesticide concentrates.

This product is not for use on medical device surfaces.

Before use in federally inspected meat and poultry food processing plants and dairies, food products and packaging materials must be removed from room or carefully protected. A potable water rinse is not allowed (Do not rinse) following use of the product as a sanitizer on previously cleaned hard surfaces provided that the surfaces are adequately drained before contact with food so that little or no residue remains.

### Limitations, Restrictions, and Exceptions

For direct injection into spray waters used on process lines

Treat water containing plant pathogens by injecting SaniDate 5.0 Sanitizer/Disinfectant directly into spray system water with 0.05 fl. oz. of concentrate for every 1 gallon of water or use a dilution rate of 1:2,500. Applicable for use on all types of post-harvest commodities.

### Method

[Directed](#)

[Injection](#)

[Spray](#)

Rates

[field\\_rates 0](#)

- 

Timings

[Post-harvest](#)