

POST-HARVEST SPRAY TREATMENTS ON PROCESS AND PACKING LINES

General Information

DIRECTIONS FOR USE

SaniDate 5.0 Sanitizer/Disinfectant works best when diluted with water containing minimal levels of organic or inorganic materials, and with water having a neutral pH. Thoroughly rinse out tank with water before mixing concentrate. This product will readily mix with clean, neutral water and does not require agitation.

SaniDate 5.0 Sanitizer/Disinfectant concentrate should not be combined or mixed with any other pesticide concentrates.

This product is not for use on medical device surfaces.

Before use in federally inspected meat and poultry food processing plants and dairies, food products and packaging materials must be removed from room or carefully protected. A potable water rinse is not allowed (Do not rinse) following use of the product as a sanitizer on previously cleaned hard surfaces provided that the surfaces are adequately drained before contact with food so that little or no residue remains.

Limitations, Restrictions, and Exceptions

For post-harvest spray treatments on process and packing lines

Inject SaniDate 5.0 Sanitizer/Disinfectant directly into spray system water on process and packing lines to prevent bacterial and fungal diseases on post-harvest fruits and vegetables. Inject at a rate of 1:250 - 1:2,500 concentrate to clean water. For best results, where dump tanks are used, make post harvest spray treatment as produce is leaving dump tanks. Applicable for use on all types of post harvest commodities.

Method

[Injection](#)

[Spray](#)

-

Timings

[Post-harvest](#)